

Chapter 2

Detection of 2-Alkyl-4-Quinolones Using Biosensors

Stephen P. Diggle, Matthew P. Fletcher, Miguel Cámara,
and Paul Williams

Abstract

2-Alkyl-4-quinolones (AQs) such as 2-heptyl-3-hydroxy-4-quinolone (PQS) and 2-heptyl-4-quinolone (HHQ) are quorum sensing signal molecules. Here we describe two methods for AQ detection and quantification that employ thin layer chromatography (TLC) and microtitre plate assays in combination with a *lux*-based *Pseudomonas aeruginosa* AQ biosensor strain. For TLC detection, organic solvent extracts of bacterial cells or spent culture supernatants are chromatographed on TLC plates, which are then dried and overlaid with the AQ biosensor. After detection by the bioreporter, AQs appear as both luminescent and green (pyocyanin) spots. For the microtitre assay, either spent bacterial culture supernatants or extracts are added to a growth medium containing the AQ biosensor. Light output by the bioreporter is proportional to the AQ content of the sample. The assays described are simple to perform, do not require sophisticated instrumentation, and are highly amenable to screening large numbers of bacterial samples.

Key words: *Pseudomonas aeruginosa*, Biosensor, 2-Alkyl-4-quinolones, 2-Heptyl-3-hydroxy-4-quinolone, 2-Heptyl-4-quinolone, *pqsA*

1. Introduction

In *Pseudomonas aeruginosa*, cell–cell communication (quorum sensing, QS) is known to control the production of extracellular virulence factors and promote biofilm maturation. The QS system consists of two *N*-acylhomoserine lactone (AHL) regulatory circuits (*las* and *rhl*) linked to a 2-alkyl-4-quinolone (AQ) system (1, 2). In the *las* system, the *lasI* gene product directs the synthesis of *N*-(3-oxo-dodecanoyl)-L-homoserine lactone (3-oxo-C₁₂-HSL), which interacts with the transcriptional regulator LasR and activates target promoters. In the *rhl* system, *rhlI* directs the synthesis of *N*-(butanoyl)-L-homoserine lactone (C₄-HSL), which

interacts with the cognate regulator RhlR and activates target gene promoters. The *las* and *rhl* systems are hierarchically connected and have been found to regulate the timing and production of multiple virulence factors including elastase, alkaline protease, exotoxin A, rhamnolipids, pyocyanin, lectins, superoxide dismutases, and biofilm formation (1).

Pesci et al. demonstrated the addition of a spent culture medium extract from a *P. aeruginosa* wild type (PAO1)-induced expression of the elastase gene *lasB* in a PAO1 AHL-deficient *lasR* mutant (3). These data suggested that a non-AHL signal produced by the bacterium was capable of activating *lasB*. It was also shown that the novel signal required a functional *rhl* system for its bioactivity, as *lasB* could not be activated in an *rhlI/rhlR* double mutant by PAO1 spent culture extracts. The molecule responsible for the non-AHL-mediated QS signalling pathway was purified and chemically identified as 2-heptyl-3-hydroxy-4-quinolone and termed the pseudomonas quinolone signal (PQS) (3) (Fig. 1). PQS belongs to the AQ family of compounds, which were first chemically identified in the 1940s and studied for their antibacterial properties. In addition to PQS, other major molecules produced by this organism that belong to this family include 2-heptyl-4-quinolone (HHQ) (Fig. 1), 2-nonyl-4-quinolone (NHQ), and 2-heptyl-4-quinolone *N*-oxide (HQNO) (4). More recently, it has been shown that AQs are produced by other genera of bacteria including *Burkholderia pseudomallei*, the causative organism of melioidosis (5), suggesting that AQ signalling may be more widespread than previously thought.

For detection of AQs, both NMR (nuclear magnetic resonance) and LC-MS (liquid chromatography-mass spectroscopy) have been used (6, 7). These methods are extremely sensitive but




Fig. 1. Structures of the AQs PQS and HHQ.

rely on access to expensive instrumentation and require considerable expertise. An alternative is to use thin layer chromatography (TLC) to detect AQs and previously TLC assays have been used to detect PQS under UV light (8–10). However, these are not discriminatory in complex bacterial supernatants and are not particularly suitable for 2-alkyl-4-quinolones such as HHQ, which fluoresces much more weakly under UV light than PQS. A third option is via the use of a specific AQ biosensor (11–13) and this chapter will describe this method of AQ detection.

2. Materials

2.1. Bacterial Bioreporters, Growth Media, and AQ Compounds

1. The bioreporter used in this assay is PAO1 $\Delta pqsA$ CTX-*lux::pqsA* (13). This strain is *P. aeruginosa* PAO1 with a chromosomal deletion in the *pqsA* gene. It is AQ negative but responds to the presence of AQs such as PQS and HHQ. The reporter gene construct in this strain is the *pqsA* promoter (which is sensitive to both PQS and HHQ) fused to a CTX-*luxCDABE* cassette. The resulting construct was conjugated into PAO1 $\Delta pqsA$ resulting in a stable chromosomal single copy reporter. The strain can be maintained in tetracycline 125 μ g/ml and stored at -80°C in 25% (vol/vol) glycerol.
2. PQS standard, 10 mM in methanol (PQS MW 259).
3. HHQ standard, 10 mM in methanol (HHQ MW 243).
4. Luria-Bertani (LB) medium: 1% (wt/vol) bacto-tryptone, 0.5% (wt/vol) yeast extract, 1% (wt/vol) sodium chloride in distilled water.
5. LB agar: LB media with addition of 2% (wt/vol) agar technical no. 3.
6. Tetracycline, 50 mg/ml in methanol.

2.2. Extraction of AQs

1. Potassium phosphate monobasic (KH_2PO_4).
2. Methanol-HPLC gradient grade.
3. Ethyl acetate-HPLC gradient grade.
4. Acetone-HPLC gradient grade.
5. Dichloromethane-HPLC gradient grade.
6. Glacial acetic acid-analytical grade.
7. Rotary evaporator (e.g. R-114, Buchi).

2.3. Detection of AQs Using TLC

1. Normal phase 20×20 cm silica 60_{F254} TLC plates (Merck).
2. TLC developing tank (Sigma-Aldrich, $27.5 \times 27.5 \times 7.5$ cm).
3. Soft top agar: 0.65% (wt/vol) agar technical no. 3, 1% (wt/vol) tryptone, 0.5% (wt/vol) sodium chloride.

4. UV transilluminator (e.g. Vilber Lourmat TFX-20.M, 312 nm).
5. Luminograph photon video camera (e.g. LB 980, EG & G Berthold).
6. X-ray film (e.g. Pierce CL-Xposure 18×24 cm film).

2.4. Detection of AQs Using Microtitre Plates

1. 96-Well plate luminometer (e.g. Anthos Lucy1 or Tecan GENios Pro).
2. 96-Well, black, clear bottom microtitre plates (e.g. Costar-Corning Inc.).

3. Methods

3.1. Preparing Bacterial Cultures for AQ Extraction

1. Under sterile conditions streak out a 10 µl loop of the test bacterium, *P. aeruginosa* PAO1 (positive control), the AQ-negative mutant PAO1Δ*pqsA* (AQ negative control), and the PQS-negative (but HHQ positive) PAO1Δ*pqsH* (PQS negative control) onto fresh LB agar plates. Grow overnight at 37°C.
2. On the following day, inoculate 5 ml of LB medium with single colonies of the relevant strains. Grow the cultures overnight at 37°C with shaking at 200 rpm. It is not essential that LB medium is used for this stage, any nutrient medium can be used, although *P. aeruginosa* strains produced high concentrations of AQs in LB. If an individual strain requires antibiotics for selection, then these can be added at this stage.
3. The following day, the optical densities (OD) of the cultures should be measured at 600 nm (OD₆₀₀). Using these readings, standardise the cultures to approximately OD 1.0 by diluting with the growth medium used in step 2.
4. Using sterile techniques, transfer 0.25 ml of standardised culture into 250 ml Erlenmeyer flasks containing 25 ml LB broth (or alternative growth medium). A small volume of culture in a large flask allows good aeration of the medium. Incubate at 37°C, with shaking at 200 rpm for 8 h (see Note 1).

3.2. AQ Extraction of Bacterial Cultures

1. Transfer a defined volume of each culture (in this example 10 ml) to a 50 ml centrifuge tube and centrifuge at 10,000×*g* for 10 min. Extract the cells (see step 2) or the supernatant (see step 3). It is possible to carry out both procedures together.
2. For extraction of AQs from cells, centrifuge the culture at 10,000×*g* for 10 min, remove the supernatant (save if required for supernatant extraction) and resuspend the cells in 10 ml of PBS buffer. Centrifuge again at 10,000×*g* for

10 min and discard the supernatant. Repeat the media wash steps twice to remove all traces of the supernatant AQs from the cells. Add 10 ml of methanol to the cell pellet and vortex until fully resuspended. Allow to stand for 10 min to allow the cells to lyse before centrifuging again at $10,000 \times g$ for 10 min. Filter the extract through sterile Minisart (Sartorius) 0.2 μM filters into clean centrifuge tubes to remove all cell debris from the extraction mixtures. At this stage, it is possible to store the cell extractions in the freezer at -20°C for several days if required.

3. For supernatant extraction, centrifuge the culture at $10,000 \times g$ for 10 min and filter the supernatants through sterile Minisart (Sartorius) 0.2 μM filters into clean centrifuge tubes to remove any cells. Add 10 ml of acidified ethyl acetate (glacial acetic acid 0.01% (vol/vol) in ethyl acetate) to the supernatant and vortex for 30 s so the two phases are well mixed. Transfer the extraction mixtures into a separating funnel that has been previously washed with acetone and allow the extraction mixtures to settle and the two phases to separate. Transfer the top organic layer into a fresh centrifuge tube. Repeat the extraction procedure on the bottom layer twice before discarding. Pool the collected organic layers. If time is limiting, supernatant extraction mixtures can be stored in the freezer at -20°C for several days.
4. For both the cell and supernatant procedures, transfer the extraction mixtures to 50 ml round bottom flasks that have been previously washed with acetone and rotary evaporate the mixtures to dryness.
5. Add 0.5 ml of methanol to the round bottom flasks and agitate for 30 s before transferring the liquid to 2 ml glass sample vials. Repeat this step with two further additions of 0.5 ml methanol and pool each sample in each vial. If time is limiting, both cell and supernatant extraction mixtures can be stored in the freezer at -20°C for several days.
6. Dry down the extraction mixtures in the sample vials under a stream of nitrogen gas. Dry cell and supernatant extraction residues can be stored in the freezer at -20°C for several months.

3.3. Preparation of TLC Plates and Running of Samples

1. Prepare normal phase silica 20×20 cm 60_{F254} TLC plates by soaking in a 5% (wt/vol) solution of KH_2PO_4 for 30 min before activating at 100°C for 1 h. A hybridization oven can be used to do this.
2. Draw a faint line in pencil approximately 1 in. from the bottom of the silica TLC plate to use as a guide for spotting sample extracts.

3. Reconstitute sample extracts in 100 µl of methanol and spot 5 µl of each onto the TLC plate. The amount spotted onto the TLC depends on the concentration of AQs in the sample. *P. aeruginosa* produces high amounts of AQs and in this case, 5 µl is a good starting quantity. As positive controls, 2 µl of 10 mM stock solutions of PQS and HHQ (or other AQs) in methanol can be spotted onto the TLC plate. Space each spot at 2 cm intervals along the line. A hairdryer can be used to dry samples during spotting to give a tighter spot.
4. When dry, place the TLC plate into a developing tank and run the TLC using a mixture of dichloromethane:methanol (95:5) as the mobile phase until the solvent front is 1–2 cm from the top of the plate. The TLC plate can be visualised using a UV transilluminator at 312 nm and photographed at this point.
5. Allow the TLC plate to dry and apply autoclave tape to both the underside of the TLC plate and around the edges so that the tape creates a well at least 0.5 cm deep around the TLC plate into which the nutrient agar containing the biosensor will be poured. Make sure the autoclave tape is firmly pressed down and forms a tight seal (Fig. 2).

3.4. Overlay of TLC Plates and Detection of AQs Using a Bioreporter

1. Streak out a 10 µl loop of the AQ biosensor (*PAO1ΔpqsA CTX-lux::pqsA*) (13) onto fresh LB agar plates containing tetracycline 125 µg/ml and grow overnight at 37°C.
2. The following day, inoculate a single colony into 5 ml LB medium containing tetracycline 125 µg/ml and grow overnight at 37°C with shaking at 200 rpm.
3. The following day, gently melt 100 ml of soft top agar in a microwave and allow to cool to ~50°C. Add 1 ml of the overnight culture to the soft top agar and mix gently (see Note 2).
4. Pour the agar mixture slowly into the well made around the TLC plate, being careful to minimise bubble formation in the agar and on the TLC plate (see Note 3).
5. Allow the agar to solidify around a Bunsen flame (to help keep sterile) and then incubate the plate at 37°C for 6 h to view light production, or overnight to view pyocyanin production. Visualise the plates for light production using a luminograph photon video camera (or similar) or develop using X-ray film (e.g. Pierce CL-XPosure film). Alternatively simply view production of the blue/green phenazine pigment pyocyanin by eye (Fig. 3) (see Note 4).

3.5. Testing for AQ Production Using a Microtitre Plate Assay

1. Prepare crude bacterial culture supernatants by growing the test bacterium as described in Subheading 3.1.
2. Remove 5 ml of culture and spin at 10,000×g for 5 min before collecting the supernatant and passing it through a




Fig. 2. Preparation of the biosensor bacteria agar TLC plate overlay. (a) Attach a strip of autoclave tape to the aluminium backing along each edge of the TLC plate, pressing down firmly to ensure a good seal. (b) Neatly trim the excess autoclave tape using scissors. (c) Pinch the autoclave tape at each corner of the TLC plate, creating a barrier of autoclave tape along each side of the TLC plate to create a well. (d) Place the TLC plate prepared as shown in (c) into an appropriate dish and pour into the well the molten soft top agar containing the biosensor bacteria.

sterile Minisart (Sartorius) 0.2 μm filter into clean tubes (see Note 5). If time is limiting, this supernatant extract can be frozen for a few days.

3. Grow the AQ biosensor overnight and dilute with LB medium to OD_{600} 1.0. Further dilute this standardised biosensor culture with LB medium to give (a) 1 in 50 and (b) 1 in 100 dilutions.
4. Sterilise a 96-well plate, e.g. under strong UV light for 15 min (see Note 6). For each well, mix 100 μl of the sterile test bacterial supernatant with 100 μl of the 1 in 50 dilutions of the biosensor to give a final culture dilution of 1 in 100. For each negative control well, add 200 μl of the 1 in 100 dilutions of the AQ biosensor. A positive control well containing a PQS or HHQ standard at a concentration of 25 μM can be added or alternatively a *P. aeruginosa* culture supernatant (100 μl) can also be included in a well during the assay.




Fig. 3. TLC assay for Aqs. (a) TLC plate run with standards of PQS and HHQ and supernatant extracts of PAO1, PAO1 $\Delta pqsH$, and PAO1 $\Delta pqsA$ and visualized under UV at 312 nm. (b) Overlay of TLC plate with soft top agar containing biosensor bacteria showing the production of light in response to Aqs visualized using a luminograph photon video camera. (c) Overlay of TLC plate with soft top agar containing the biosensor bacteria showing production of the blue/green pigment, pyocyanin, in response to Aqs. TLC lanes: (1) PQS 10 mM, 2 μ l, (2) HHQ 10 mM, 2 μ l, (3) PAO1 supernatant extract, 5 μ l, (4) PAO1 $\Delta pqsH$ supernatant extract, 5 μ l, (5) PAO1 $\Delta pqsA$ supernatant extract, 5 μ l. The AQ biosensor emits light over spots identified as PQS and HHQ in PAO1 and HHQ only in PAO1 $\Delta pqsH$. No light spots are observed for the AQ-negative *pqsA* mutant.

5. Monitor bioluminescence and OD at 37°C using a combined spectrophotometer/luminometer, e.g. the Anthos LUCY1 controlled by the Stingray software (Dazdaq). This measures OD and bioluminescence from all wells every 30 min for 24 h. Luminescence is recorded as relative light units (RLU)

per unit of OD. If an automated combined spectrophotometer/luminometer is unavailable, readings can be taken manually at defined time points by growing the bacterial cultures under specific conditions and measuring the OD and bioluminescence of culture samples using a spectrophotometer and tube luminometer (e.g. EG & G Junior), respectively.

4. Notes

1. Growth of *P. aeruginosa* cultures for 8 h in LB medium results in the bacteria reaching mid-stationary phase, which is sufficient for high quantities of AQs to be produced. If using an alternative growth medium, the time of incubation may have to be altered to accommodate the bacterial cells reaching the stationary phase of growth. We recommend that stationary phase cells be used when attempting to detect AQs produced by bacterial species.
2. Make sure the agar has cooled sufficiently before adding the bacterial reporter strain. Addition at too high a temperature will attenuate bacterial growth.
3. Pour the agar promptly or it will begin to solidify. Bubbles can be removed by gently passing a Bunsen burner flame over the surface of the agar.
4. Both PQS and HHQ will activate light production in the reporter, as both control the expression of the *pqsA* gene. In addition, both PQS and HHQ are required for the production of pyocyanin.
5. In addition to cell-free culture supernatants, the solvent extracted culture extracts described in Subheading 3.2 may also be analysed via this method. Simply dilute 5 µl of the solvent extract in 100 µl of LB and add to 100 µl of 1 in 50 dilutions of the AQ biosensor per well.
6. Specialised 96-well plates need to be used when monitoring bioluminescence. The plates need to have black sides to reduce light scatter between wells and have clear plastic bottoms so that an automated luminometer can detect and measure light output accurately.

Acknowledgments

We gratefully acknowledge the Royal Society (SPD) and BBSRC (MPF) for funding.

References

1. Williams, P., and Cámara, M. (2009) Quorum sensing and environmental adaptation in *Pseudomonas aeruginosa*: a tale of regulatory networks and multifunctional signal molecules. *Curr. Opin. Microbiol.* **12**, 182–191.
2. Popat, R., Crusz, S. A., and Diggle, S. P. (2008) The social behaviours of bacterial pathogens. *Br. Med. Bull.* **87**, 63–75.
3. Pesci, E. C., Milbank, J. B., Pearson, J. P., McKnight, S., Kende, A. S., et al. (1999) Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA.* **96**, 11229–11234.
4. Dubern, J. F., and Diggle, S. P. (2008) Quorum sensing by 2-alkyl-4-quinolones in *Pseudomonas aeruginosa* and other bacterial species. *Mol. Biosyst.* **4**, 882–888.
5. Diggle, S. P., Lumjiaktase, P., Dipilato, F., Winzer, K., Kunakorn, M., et al. (2006) Functional genetic analysis reveals a 2-alkyl-4-quinolone signaling system in the human pathogen *Burkholderia pseudomallei* and related bacteria. *Chem. Biol.* **13**, 701–710.
6. Lépine, F., Déziel, E., Milot, S., and Rahme, L. G. (2003) A stable isotope dilution assay for the quantification of the *Pseudomonas aeruginosa* quinolone signal in *Pseudomonas aeruginosa* cultures. *Biochim. Biophys. Acta.* **1622**, 36–41.
7. Lépine, F., Milot, S., Déziel, E., He, J., and Rahme, L. G. (2004) Electrospray/mass spectrometric identification and analysis of 4-hydroxy-2-alkylquinolines (HAQs) produced by *Pseudomonas aeruginosa*. *J. Am. Soc. Mass Spectrom.* **15**, 862–869.
8. Diggle, S. P., Winzer, K., Chhabra, S. R., Worrall, K. E., Cámara, M., and Williams, P. (2003) The *Pseudomonas aeruginosa* quinolone signal molecule overcomes the cell density-dependency of the quorum sensing hierarchy, regulates *rhl*-dependent genes at the onset of stationary phase and can be produced in the absence of LasR. *Mol. Microbiol.* **50**, 29–43.
9. Calfee, M. W., Coleman, J. P., and Pesci, E. C. (2001) Interference with *Pseudomonas aeruginosa* quinolone signal synthesis inhibits virulence factor expression by *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA.* **98**, 11633–11637.
10. Gallagher, L. A., McKnight, S. L., Kuznetsova, M. S., Pesci, E. C., and Manoil, C. (2002) Functions required for extracellular quinolone signaling by *Pseudomonas aeruginosa*. *J. Bacteriol.* **184**, 6472–6480.
11. Fletcher, M. P., Diggle, S. P., Crusz, S. A., Chhabra, S. R., Cámara, M., and Williams, P. (2007) A dual biosensor for 2-alkyl-4-quinolone quorum-sensing signal molecules. *Environ. Microbiol.* **9**, 2683–2693.
12. Fletcher, M. P., Diggle, S. P., Cámara, M., and Williams, P. (2007) Biosensor-based assays for PQS, HHQ and related 2-alkyl-4-quinolone quorum sensing signal molecules. *Nat. Protoc.* **2**, 1254–1262.
13. Diggle, S. P., Matthijs, S., Wright, V.J., Fletcher, M. P., Chhabra, S. R., et al. (2007) The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS play multifunctional roles in quorum sensing and iron entrapment. *Chem. Biol.* **14**, 87–96.