

Residual sarcoplasmic reticulum Ca^{2+} concentration after Ca^{2+} release in skeletal myofibers from young adult and old mice

Zhong-Min Wang · Shen Tang · María Laura Messi · Jenny J. Yang · Osvaldo Delbono

Received: 28 November 2011 / Revised: 31 December 2011 / Accepted: 2 January 2012 / Published online: 17 January 2012
© Springer-Verlag 2012

Abstract Contrasting information suggests either almost complete depletion of sarcoplasmic reticulum (SR) Ca^{2+} or significant residual Ca^{2+} concentration after prolonged depolarization of the skeletal muscle fiber. The primary obstacle to resolving this controversy is the lack of genetically encoded Ca^{2+} indicators targeted to the SR that exhibit low- Ca^{2+} affinity, a fast biosensor: Ca^{2+} off-rate reaction, and can be expressed in myofibers from adult and older adult mammalian species. This work used the recently designed low-affinity Ca^{2+} sensor ($K_d=1.66$ mM in the myofiber) CatchER (calcium sensor for detecting high concentrations in the ER) targeted to the SR, to investigate whether prolonged skeletal muscle fiber depolarization significantly alters residual SR Ca^{2+} with aging. We found CatchER a proper tool to investigate SR Ca^{2+} depletion in young adult and older adult mice, consistently tracking SR luminal Ca^{2+} release in response to brief and repetitive stimulation. We evoked SR Ca^{2+} release in whole-cell voltage-clamped flexor digitorum brevis muscle fibers from young and old FVB mice and tested the maximal SR Ca^{2+} release by directly activating the ryanodine receptor (RyR1) with 4-chloro-*m*-cresol in the same myofibers. Here, we report for the first time that the Ca^{2+} remaining in the SR after prolonged depolarization (2 s) in myofibers from aging (~220 μM) was larger than young (~132 μM) mice. These

experiments indicate that SR Ca^{2+} is far from fully depleted under physiological conditions throughout life, and support the concept of excitation–contraction uncoupling in functional senescent myofibers.

Keywords Skeletal muscle · Calcium sensor · CatchER · Sarcoplasmic reticulum · Aging

Abbreviations

ER/SR	Endoplasmic/sarcoplasmic reticulum
EGFP	Enhanced green fluorescence protein
FDB	Flexor digitorum brevis
RyR1	Ryanodine receptor-isoform-1
4-CmC	4-Chloro- <i>m</i> -cresol
TEA	Tetraethylammonium
di-8-	di-8-amino naphthyl ethenyl pyridinium
ANEPPS	
CPA	Cyclopiazonic acid
CatchER	Calcium sensor for detecting high concentrations in the ER
GECI	Genetically encoded Ca^{2+} biosensor

Introduction

Measuring SR residual Ca^{2+} concentration in depolarized skeletal muscle fiber is essential to understanding the influence of luminal SR Ca^{2+} on SR Ca^{2+} release and force deterioration with aging. Developing reliable molecular tools to detect Ca^{2+} kinetics in intracellular organelles is imperative to determining its role in cell signaling, physiological responses to training and exercise, and the pathogenesis of muscular dystrophy, central core disease, malignant hyperthermia, and fatigue syndromes [27, 44]. The SR is the

Z.-M. Wang · M. L. Messi · O. Delbono (✉)
Department of Internal Medicine, Section on Gerontology
and Geriatric Medicine,
Wake Forest University School of Medicine,
1 Medical Center Boulevard,
Winston-Salem, NC 27157, USA
e-mail: odelbono@wfubmc.edu

S. Tang · J. J. Yang
Department of Chemistry, Center
for Drug Design and Biotechnology, Georgia State University,
Atlanta, GA 30303, USA

main source of Ca^{2+} for skeletal muscle contraction, but the role of its intricate terminal cisternae/longitudinal architecture and Ca^{2+} buffering by endogenous Ca^{2+} -binding proteins, such as calsequestrin and calreticulin, in intracellular Ca^{2+} homeostasis is poorly understood. Particularly, the elevated SR Ca^{2+} concentration requires a low-affinity biosensor to reliably monitor Ca^{2+} concentration and kinetics.

Modifications in Ca^{2+} concentration are the signal that regulates a number of functions in excitable and nonexcitable cells [15, 16, 36]. However, loading the skeletal muscle SR with Ca^{2+} -sensitive indicators presents technical difficulties that seriously hamper our ability to determine Ca^{2+} concentration and release kinetics and to answer questions relevant to muscle physiology and pathology. Assessing residual SR Ca^{2+} in excitable cells is relevant for the study of luminal Ca^{2+} buffer capacity, Ca^{2+} -dependent regulation of IP3R and RyR function, store-operated Ca^{2+} entry, termination of SR Ca^{2+} release, and age-dependent decline in SR Ca^{2+} release [12]. Therefore, measuring Ca^{2+} movements using engineered protein targeted to the endoplasmic/sarcoplasmic reticulum (ER/SR) is increasingly important [4, 12, 19, 28, 34].

Previous studies have shown that decreased expression of the $\text{Ca}_v1.1$ subunit diminishes SR Ca^{2+} release in muscle fibers from aging mice [11, 12], a process known as excitation–contraction uncoupling (ECU) [3]. ECU is partially due to increased $\text{Ca}_v\beta1a$ expression with aging [35]. Recently, we estimated SR Ca^{2+} depletion in response to 4-chloro-*m*-cresol (4-CmC) or myofiber depolarization in myofibers from young adult mice electroporated with the Ca^{2+} biosensor D1ER [12]. We concluded that significant depletion of SR Ca^{2+} under physiological conditions is unlikely in young mice [12]. However, quantifying SR Ca^{2+} concentration after prolonged myofiber depolarization was not possible due to limitations in calibrating D1ER in the myofiber as described previously [28]. Quantifying residual SR Ca^{2+} requires measuring its resting concentration, release, and content in the same fiber using CatchER as the Ca^{2+} biosensor.

We recently reported that the novel enhanced green fluorescence protein-derived biosensor CatchER can monitor Ca^{2+} concentration and dynamics in high Ca^{2+} concentration compartments, such as the SR. It exhibits relatively low affinity in the skeletal muscle, as its K_d , calibrated in the myofiber, is 1.66 mM, and an unprecedented off-rate at $\sim 700 \text{ s}^{-1}$, which makes it one of the more advanced and useful tools for examining changes in SR Ca^{2+} concentration in response to pulses both as prolonged as those applied to elicit muscle tetanus and as short as an action potential [34].

Here, we show that CatchER exhibits a consistent response to muscle fiber stimulation, which allows us to quantify SR Ca^{2+} content, the rate of Ca^{2+} flux in the SR lumen, and the residual SR Ca^{2+} concentration in response to prolonged electrical stimulation beyond physiological demands in

voltage-clamped adult mouse flexor digitorum brevis (FDB) muscle fibers. Maximal SR Ca^{2+} depletion was also examined by bypassing the physiological $\text{Ca}_v1.1$ /ryanodine receptor-1 (RyR1) coupling to stimulate the RyR1 directly. For this application, we used the specific RyR1 agonist 4-chloro-*m*-cresol based on the reproducibility of its action and the lack of any reported effect on SR Ca^{2+} refilling.

We report a significant larger residual SR Ca^{2+} concentration in fibers from old compared to young adult mice, finding that supports the concept of excitation–contraction uncoupling in senescent mice.

Materials and methods

Flexor digitorum brevis muscle fibers

Single skeletal muscle fibers from the FDB were obtained from 3- to 5-month-old C57BL6 mice raised in the Wake Forest University School of Medicine (WFUSM) Animal Research Program. Mice were killed by cervical dislocation following isoflurane anesthesia. For aging studies, we used young (3–5 months) and old (22–25 months) FVB (*Freund virus B*) mice from the WFUSM colony. Animal handling followed a protocol approved by the WFUSM Animal Care and Use Committee.

FDB fiber electroporation and SR Ca^{2+} fluorescence recording

The pcDNA3.1 plasmid carrying the CatchER biosensor was electroporated into the FDB muscle as described [6]. FDB fibers were enzymatically dissociated and recorded 2–3 weeks after electroporation [34]. Dissociated FDB fibers were voltage-clamped in the whole-cell configuration of the patch-clamp technique [9], according to procedures specific for muscle fibers [39]. Potential voltage-errors associated with whole-cell recording in large cells were minimized by selecting small FDB fibers and adequately compensating for whole-cell capacitance transients. The pipette electrode was filled with the following solution (mM): 130 Cs-aspartate, 2 MgCl_2 , 0.2 or 20 Cs_2EGTA as specified below, 10 HEPES, 5 Na-ATP, and 0.5 GTP; pH was adjusted to 7.3 with CsOH. High EGTA was used to reproduce experimental conditions in a series of studies on FDB muscle fibers in which SR Ca^{2+} release was recorded in the cytosol [12, 34, 41]. The electrode resistance ranged from 450 to 650 k Ω . The external solution contained (in mM): 100 TEA-OH (tetraethylammonium hydroxide), 50 Na_2SO_4 , 2 MgSO_4 , 2 CaSO_4 , 2 3-4DAP, and 5 Na-HEPES. Solution pH was adjusted to 7.3 with CH_3SO_3 [12, 39]. These solutions ensured good control of resting leakage currents, and the preparation was preserved throughout the experiment. Methanesulphonic acid is the counterion to TEA. Drift and

noise imposed by silver-chloride wires were corrected by standard methods. All experiments were recorded at room temperature (21–22°C). To permeate the myofibers, we used a solution containing 90 mM K-glutamate, 1.02 mM MgCl₂, 5 mM NaCl, 10 mM HEPES, 1 mM 1,2-bis(*o*-aminophenoxy)ethane-*n,n,n',n'*-tetraacetic acid (BAPTA), 0.323 mM CaCl₂, 0.025 *N*-benzyl-*p*-toluene sulphonamide, 2% (vol/vol) poly (*N*-vinyl-2-pyrrolidone) (1 mM free Mg²⁺, 0.0001 mM free Ca²⁺), pH 7.2, adjusted with KOH [43]. Free [Ca²⁺] and [Mg²⁺] in solution were calculated by the Max-Chelator program.

Fluorescence was recorded using an Axiovert 200 microscope with a 20×/0.75 (Zeiss, Oberkochen, Germany) and a Radiance 2100 (Bio-Rad, Zeiss) confocal system. Fibers were imaged through a C-Apochromat 40× water-immersion objective (NA 1.2, Zeiss) or a 20× Fluor (NA 0.75) using a krypton–argon laser at 488-nm excitation wavelength. CatchER's fluorescence emission was measured at 528±25 nm wavelength. For most experiments, the laser was attenuated to 6–12% with a neutral density filter. Fibers were imaged in line-scan (*x*–*t*) mode. The fiber was always oriented parallel to the *x* scan direction. Line-scan images were acquired with 256 pixels (0.236 mm/pixel) in the *x*- and 512 pixels (0.833 ms/pixel) in the *t*-direction. For image acquisition, we used LaserSharp 2000 software (Bio-Rad, Zeiss). Cytosolic Ca²⁺ transients were recorded using 5 μM Rhod-2 AM (Invitrogen, Carlsbad, CA) loaded for 30 min on CatchER-expressing FDB fibers. Fibers were excited at 568 nm using a krypton laser and recorded at 600 nm.

SR Ca²⁺ release and cytosolic Ca²⁺ transients were recorded sequentially (20 s interval). No differences in the amplitude and kinetics of SR Ca²⁺ release monitored with CatchER were apparent in Rhod-2-loaded and unloaded fibers (data not shown).

CatchER calibration in mouse FDB muscle fibers was reported in detail previously (Supporting Information in [34]).

Statistics

Values are given as mean±SEM with the number of observations (*n*). Statistical analysis was performed using the Student's unpaired *t* test and the Mann–Whitney rank-sum test when values were not normally distributed. *P*<0.05 was considered significant.

Results

RyR1 mediates SR Ca²⁺ release recorded with CatchER

To verify that RyR1 mediates the Ca²⁺ release detected with CatchER, we exposed fibers to the channel blocker

ryanodine. Nadir SR Ca²⁺ fluorescence decreased with time after applying 5 μM ryanodine extracellularly. Figure 1 illustrates CatchER fluorescence in response to a series of 20 mV/100 ms pulses delivered before (*a*) and after (*b*, *c*) applying ryanodine to fibers from young adult mice. The amplitude of CatchER fluorescence 28 and 55 min after applying the drug compared to the amplitude of the signal at time zero was 45±4% and 0.3±0.02% (*n*=3). During ryanodine application, basal ΔF/F did not change significantly; it was 1.61±0.27 and 1.49±0.28 at times 0 and 55 min, respectively. Since RyR1 is the only RyR isoform expressed in adult skeletal muscle, we conclude that it mediates decreases in SR Ca²⁺ in response to sarcolemmal depolarization [2, 22, 26]. Declines in CatchER fluorescence, in response to the agonist 4-CmC, support this conclusion (see below). The slow time scale of the effect indicates slow diffusion of the agent. The kinetics of ryanodine binding to its receptor may be a contributing factor, as reported [12]. Control experiments in which ryanodine was omitted exhibited a maximal decline of 20±2.3% (*n*=5) in CatchER's ΔF/F at the end of 60 min.

High cytosolic EGTA concentration does not affect SR Ca²⁺ recovery

Like D1ER [12], CatchER displays a fluorescence signal larger in the presence of 20 mM than in 0.2 mM cytosolic EGTA; despite EGTA's Ca²⁺ buffer capacity, SR Ca²⁺ recovery is not altered. Figure 2a shows the time course of CatchER fluorescence in response to 20 mV/100 ms command pulses recorded in 0.2 (black trace) and 20 mM (red trace) EGTA in fibers from young adult mice. CatchER signal in 0.2 mM EGTA was scaled up to overlap with its




Fig. 1 RyR1 mediates SR Ca²⁺ release. Time course of CatchER ΔF/F in control (first 5 min) and after adding 5 μM ryanodine to the bath solution. Traces *a*, *b*, and *c* represent SR Ca²⁺ transients monitored with CatchER in control, 33, and 60 min, respectively, in 3 experiments with a representative fiber




Fig. 2 High cytosolic EGTA concentration increases CatchER transient amplitude but does not affect SR Ca^{2+} recovery. **a** SR Ca^{2+} transients recorded in electroporated FDB fibers and recorded in whole-cell patch-clamp using 0.2 (black trace) or 20 mM (red trace) EGTA in the pipette solution. **b** CatchER $\Delta F/F$ –[EGTA] relationship. CatchER fluorescence at nadir of the response (**b**) is significantly larger in 20 than in 0.2 mM EGTA ($p < 0.001$, $n = 11$ in 0.2 mM EGTA; $n = 13$ in 20 mM EGTA). **c** SR Ca^{2+} recovery is expressed as $(F_c - F_b)/(F_a - F_b)$, where a is the average of the basal fluorescence points, and b and c are CatchER fluorescence at nadir and 2 s after repolarization starts, respectively. SR Ca^{2+} recovery is independent of cytosolic [EGTA] ($p > 0.05$)

recording in 20 mM EGTA to compare their kinetics. High cytosolic EGTA concentration (20 mM) does not seem to prevent SR Ca^{2+} recovery and elicits average fluorescence signals 2.6 times larger than those recorded in 0.2 mM EGTA, which is statistically significant (Fig. 2b). Why SR Ca^{2+} depletion is more pronounced with high EGTA concentration is not obvious. Perhaps in the low EGTA solution, Ca^{2+} inactivates RyR1. To examine whether strong cytosolic Ca^{2+} chelation modifies SR Ca^{2+} uptake, we analyzed SR Ca^{2+} recovery 2.5 s after the end of the depolarizing pulse. Figure 2c shows that SR Ca^{2+} recovery in 0.2 and 20 mM EGTA did not differ significantly. In addition, 38 fibers recorded with 20 mM EGTA responded to electrical stimulation, while two out of 19 in 0.2 mM EGTA did not. Differences in Ca^{2+} recovery beyond c in the two experimental conditions were not statistically significant. Based on these results, we used a pipette solution containing 20 mM EGTA for the remaining experiments.


Brief and prolonged SR Ca^{2+} release tracked with CatchER

Previously reported Ca^{2+} biosensors targeted to the SR have been used to track slow changes in luminal SR Ca^{2+} [1, 12, 28, 32, 33, 42, 44]. We demonstrated that CatchER detected SR Ca^{2+} in response to a single action potential [44]. Here, we examined whether it could detect a variety of pulses from brief repetitive to a single prolonged stimulation under voltage-clamp. Figure 3 shows changes in CatchER's fluorescence in response to various command pulses to 20 mV. These data indicate that CatchER responds adequately to pulses of different duration (a, 10 ms; e, 100 ms), frequency (b, 10; c, 3.3; d, 1.6; and f, 5 Hz), or duration and frequency (g, 50 ms pulse at 3.3 Hz) and can thus monitor SR Ca^{2+} dynamics under a variety of physiological conditions. Figure 3 shows that SR Ca^{2+} is incompletely recovered within the recording time, regardless of the pulse applied to the cell. We rule out cell damage as a potential mechanism because all myofibers included in this study maintained membrane electrical properties within 10% of the original values throughout the experiment. Photobleaching partially explains the slow fluorescence recovery. Figure 3h shows a typical change in CatchER fluorescence in response to a 30 mV/100 ms pulse (raw trace). Correcting for photobleaching (corrected trace) partially speeds recovery ($21 \pm 3.2\%$; $n = 17$ fibers), or perhaps calsequestrin binds Ca^{2+} with higher affinity than CatchER [10], delaying recovery of the biosensor's fluorescence. We also cannot rule out the influence of the CatchER: Ca^{2+} on-rate reaction.

Voltage-dependent CatchER luminal signal and cytoplasmic Ca^{2+} transients

Figure 4 shows young adult mouse FDB fibers expressing CatchER loaded with 5 μM Rhod2AM and depolarized with 100 (A, B) or 500 (C, D) ms command pulses from $V_h = -80$ mV to 20 mV applied in the whole-cell configuration of patch-clamp. Panels A and C correspond to Rhod2 while B and D to CatchER. Top panels (a) show images in line-scan mode with vertical and horizontal axes corresponding to space and time, respectively. Fluorescence increases (A, C) or decreases (B, D) depend on whether cytosolic Rhod-2 or SR CatchER fluorescence was recorded, respectively. The use of 20 mM EGTA in the internal (pipette) solution provided a dual benefit: it prevented Rhod-2 saturation and the movement of artifacts during fiber depolarization. Middle panels (b) display time-dependent changes in the Ca^{2+} indicator's fluorescence. The decreased fluorescence signal indicates that Ca^{2+} dissociates from CatchER upon fiber depolarization. The rate of fluorescence signal rise for CatchER and Rhod-2 is similar, while the cytosolic fluorescence decay is faster than the luminal Ca^{2+} recovery. The rising phase of the Rhod-2 signal was fitted to a double-exponential function with time constants of

Fig. 3 CatchER tracks SR Ca^{2+} release in response to single or repetitive prolonged stimulation. Transient changes in CatchER's fluorescence response to various 20-mV command pulses: **a** a 10-ms pulse, **b** five 10-ms pulses at 10 Hz, **c** a 10-ms pulse at 3.3 Hz, **d** a 10-ms pulse at 1.6 Hz, **e**, a 100-ms pulse, **f** 100-ms pulse at 5 Hz, and **g** 50-ms pulse at 3.3 Hz. **h** CatchER's fluorescence response to a 30 mV/100 ms pulse (*raw trace, black*) and after correcting for photobleaching (*corrected trace, red*)



2 ± 0.11 and 139 ± 15 ms and 5 ± 0.31 and 246 ± 39 ms in response to 100 and 500 ms pulses, respectively. The decay phase also shows two time constants of 103 ± 17 and $1,132 \pm 168$ ms and 163 ± 19 and $1,193 \pm 145$ ($n=7$), respectively. The CatchER rising and recovery phases were fitted with a single time constant of 26 ± 3.5 and 19 ± 2.3 ms and $1,315 \pm 147$ and $2,512 \pm 323$ ($n=12$) in response to 100 and 500 ms pulses, respectively. Differences in Ca^{2+} affinity and Ca^{2+} buffer capacity of SR and cytoplasm can explain differences in the time course of CatchER and Rhod-2 fluorescence. Figure 4 (c) illustrates the Ca^{2+} currents in real-time to compare the time course of the electrical and optical signals. The time-to-peak (in ms) of the Rhod-2 signal were 98 ± 4 and 113 ± 8 , while for CatchER, the values were 91 ± 5 and 139 ± 23 in response to 100 and 500 ms pulses, respectively. Values are mean $\pm$ SEM for 14

fibers expressing CatchER and loaded with Rhod-2. Apparent differences between time-to-peak of Rhod-2 and CatchER signals were not statistically significant.

CatchER's fluorescence amplitude increases with the duration of sarcolemmal depolarization

We recently showed that CatchER's fluorescence decreases with sarcolemmal depolarization, and the fluorescence–membrane voltage relationship follows a Boltzmann equation in fibers from young and old mice [34]. To further determine whether CatchER responds to SR Ca^{2+} release as a function of pulse duration, we voltage-clamped 12 fibers from young adult mice at -80 mV and stimulated them with 20 mV command pulses of increasing duration (10, 25, 50, and 100 ms; Fig. 5a, b). As shown in Fig. 5c, CatchER's




Fig. 4 SR and cytosolic Ca^{2+} transients recorded in the same FDB fibers. Ca^{2+} transients were recorded in response to a command pulse (20 mV and 100-ms [A, B] or 500-ms [C, D] duration [$V_h = -80$ mV]). **a** shows images in confocal line-scan mode with vertical and horizontal axes corresponding to space and time, respectively; **b** Ca^{2+} transients' fluorescence profile measured with Rhod-2 (**a**, **c**) or CatchER (**b**, **d**); **c** Ca^{2+} currents. Dashed and dotted lines indicate basal fluorescence before pulse application and isoelectric current, respectively. $\Delta F/F$ for **a**, **b**, **c**, and **d** were 6.65, 0.26, 6.7, and 0.28, respectively. The rising phase of the Rhod-2 signal was fitted to an exponential function with two time constants of 3 and 146 ms and 4 and 246 ms in response to 100 and 500 ms pulses, respectively. The decay phase also shows two time constants of 114 and 1,025 ms and 159 and 1,246 ms, respectively. Time to peak is 98 and 111 ms, respectively. The CatchER rising and recovery phases were fitted with a single time constant of 31 and 23 ms and 1,214 and 2,733 ms in response to 100 and 500 ms pulses, respectively. Time to peak is 100 and 127 ms, respectively

fluorescence elicited by progressively longer pulses showed a graded increase in signal amplitude, reaching a plateau for pulses longer than 100 ms (250 and 500 ms).

Ca^{2+} remaining in the SR after prolonged depolarization differs in myofibers from young and old mice

Figure 6a shows SR Ca^{2+} fluorescence in response to a 20 mV/100 ms pulse as depicted on the top trace in a fiber from a young adult mouse. After ~2 min, the same fiber was exposed to 1 mM 4-CmC for the whole recording period, as indicated on the top bar, to bypass the excitation-SR Ca^{2+}




Fig. 5 CatchER's peak Ca^{2+} transient depends on pulse duration. **a** Various pulse durations (10, 25, 50, and 100 ms) tested in FDB fibers, voltage-clamped at -80 mV (holding potential), and pulsed to 20 mV. The pulse waveform is depicted in the top panel. **b** Maximal CatchER transient relationship to pulse duration. Data points represent the mean $\pm$ SEM of 12 fibers

release process and elicit SR Ca^{2+} release by acting directly on RyR1 (protocol 1). CatchER fluorescence seems to reach a deeper nadir than the response to the electrical pulse. The ratio between the peak response to electrical stimulation (**b**) and 4-CmC (**c**) was 0.44 ± 0.09 ($n=6$).

To investigate whether a remaining Ca^{2+} pool persists in the SR after 4-CmC application, electrical stimulation was prolonged to 500 ms, and 15 μM cyclopiazonic acid were added with 1 mM 4-CmC to block SERCA-mediated SR Ca^{2+} uptake in addition to agonist-dependent SR Ca^{2+} release (protocol 2). Increased pulse duration and the combination of the two pharmacological agents resulted in a similar SR Ca^{2+} depletion (data not shown).

Similarly, Fig. 6b shows SR Ca^{2+} fluorescence in response to a 20 mV/100 ms pulse in a fiber from an old mouse. While CatchER's response to the electrical pulse is smaller, its response to 4-CmC does not differ significantly from that obtained in young fibers [34]. The ratio between the nadir in response to electrical stimulation (**b**) and 4-CmC (**c**) was 0.21 ± 0.07 ($n=7$).

Figure 6c shows the response of myofibers from young (left) and old (right) mice to a 2 s/20 mV train of 3 ms pulses at 150 Hz. The b/c ratio was 0.41 ± 0.08 ($n=15$ fibers) and




Fig. 6 Maximal SR Ca^{2+} depletion evoked by 4-CmC. **a** CatchER fluorescence transient (bottom trace) in response to 20 mV/100 ms command pulse (top trace) in a voltage-clamped myofiber from a young adult mouse. The same fiber was exposed to 1 mM 4-CmC with ~2-min interval. The dashed line indicates the basal fluorescence (**a**), while the nadir of the response to electrical stimulation or 4-CmC application is shown in (**b**) and (**c**), respectively. **b** CatchER fluorescence transient in response to a 20 mV/100 ms command pulse in a voltage-clamped myofiber from an old mouse. The same fiber was exposed to 1 mM 4-CmC at ~2-min intervals. **c** CatchER fluorescence (bottom) in response to repetitive stimulation with 2-ms pulses at 150 Hz for 2 s at 20 mV (top) in myofibers from young (left) and old (right) mice

0.24 ± 0.06 ($n=17$ fibers), respectively. These experiments indicate that a Ca^{2+} pool remains in the SR lumen after myofiber depolarization through prolonged activation of the excitation-SR Ca^{2+} release mechanism, consistent with declining voltage-gated SR Ca^{2+} release with aging [11].

To further examine the residual luminal Ca^{2+} in the SR after prolonged Ca^{2+} release, we applied an alternative technique. After SR Ca^{2+} release was measured FDB fibers from young and old mice expressing CatchER were exposed to 0.01% saponin for 2 min in permeabilization solution (see “Materials and methods”). The permeabilized myofiber was exposed to 10^{-6} M ionomycin diluted in the previous solution to equilibrate among the extracellular space, cytosol, and lumen of the SR. Free $[\text{Mg}^{2+}]$ was set at 1 mM, and free $[\text{Ca}^{2+}]$ was set at 10^{-7} mM with BAPTA. Although apparently more stringent, this technique did not promote a significant difference in residual Ca^{2+} compared to the 4-CmC

protocols. The Ca^{2+} concentration remaining in the SR after Ca^{2+} release in response to prolonged repetitive electrical stimulation (2 s) was (in μM): 132 ± 29 and 220 ± 37 in myofibers from young ($n=9$ fibers) and old ($n=7$ fibers) mice, respectively. As resting SR Ca^{2+} concentration did not differ in myofibers from young and old mice [34], our data are consistent with a larger SR luminal Ca^{2+} depletion in fibers from young mice or an impaired voltage-gated SR Ca^{2+} release in aging mice [11].

Discussion

The main findings of this work are: (1) CatchER can track changes in SR Ca^{2+} lumen in response to pulses varying in amplitude, duration, and frequency; (2) photobleaching partially accounts for CatchER’s slow fluorescence recovery; (3) SR Ca^{2+} release monitored with CatchER is solely mediated by RyR1; (4) SR luminal Ca^{2+} release measured with CatchER mirrors the cytosolic Ca^{2+} transient, measured with Rhod-2; and (5) the fraction of residual SR Ca^{2+} is larger in myofibers from old than young mice, supporting excitation–contraction uncoupling in senescent myofibers.

CatchER is a low-affinity Ca^{2+} biosensor suitable for measuring luminal SR Ca^{2+}

CatchER allowed us to monitor changes in Ca^{2+} concentration dynamics in a practically unexplored environment—skeletal muscle SR. We recently described the structure and optical properties of these genetically encoded Ca^{2+} indicators (GECI) and the mechanism of its response to Ca^{2+} [34]. Previous attempts to measure SR Ca^{2+} in amphibian and mammalian species have been discussed elsewhere [12]. Although CatchER’s single-excitation/single-emission optical spectrum is limited compared to the single-excitation/dual emission D1ER [12, 18, 28], it has a number of advantages, including responsiveness, reproducibility of the signal, and low Ca^{2+} affinity. These properties allowed us to measure high SR luminal Ca^{2+} concentrations and their dynamics in a considerable number of myofibers after FDB muscle in vivo electroporation. Difficulties in reliably calibrating D1ER, the most commonly used biosensor for Ca^{2+} analysis in the muscle, have been reported [12, 28]; however, its ratiometric signal recorded in muscle fiber provided useful information about relative SR Ca^{2+} changes [12]. Relative changes in citrine/CFP ratio, rather than SR Ca^{2+} concentrations, have been reported using D1ER in arterial smooth muscle [42], HEK-293 cells expressing RyR2 [13], and vascular endothelial cells [21]. Here, we were able to quantify residual SR Ca^{2+} concentration after SR Ca^{2+} released in response to fiber electrical and pharmacological stimulation in young and old mice, which was

impractical with previous GECl, representing a net improvement over previous approaches. Also, CatchER expression by electroporation and recording without previous prolonged loading procedures make it a promising tool to investigate the role of SR Ca^{2+} movements under a variety of physiological and disease conditions.

The concentration of residual SR Ca^{2+} after Ca^{2+} release in skeletal myofibers from senescent mice is unknown

A series of reports estimated SR Ca^{2+} content and depletion using cytosolic synthetic Ca^{2+} indicators in young rodents. In voltage-controlled SR Ca^{2+} release in mouse FDB fibers, a 100-ms voltage pulse that maximally activates Ca^{2+} release, according to the authors, reduces the initial SR content about 80% [37]. Note that they estimated an SR Ca^{2+} content of 3 mM, consistent with previous reports [8] [7, 17, 20, 24, 30]. However, more recent works using engineered Ca^{2+} biosensors or Fluo-5N targeted to the SR showed values in the micromolar range [28, 34, 43]. We have shown that SR Ca^{2+} depletion is unlikely, even in response to prolonged depolarization using the biosensor D1ER, which was later confirmed using D4cpv [19, 32]. Using CatchER, we recorded a resting SR Ca^{2+} concentration of 512 and 573 μM in myofibers from young and old mice, respectively [34]. Some studies have reported an SR Ca^{2+} concentration of ~ 1 mM in frog muscle fibers [31, 38], while a three- to fourfold lower value (~ 308 μM) was noted using the Ca^{2+} biosensor D1ER [28]. We found higher basal SR Ca^{2+} concentration than that reported in mouse tibialis anterior muscle [28]. This difference may be explained by difficulties in measuring D1ER's R_{max} [28] or saturation due to multiple binding sites, as its reported dissociation constant is ~ 60 μM [18].

Indirect observations suggest that SR Ca^{2+} depletion does not limit its release in muscle fibers from aging mice [11]; however, no direct measure of residual Ca^{2+} content after prolonged depolarization was reported. That SR Ca^{2+} release decreases with aging would account for increased residual SR Ca^{2+} . Measurements of the residual $[\text{Ca}^{2+}]$ /resting $[\text{Ca}^{2+}]$ relationship required the development of a low-affinity biosensor targeted to the SR. Calibrations in FDB fiber showed that CatchER's K_d is 1.66 mM, which make it an excellent instrument to address this question.

Recently, SR Ca^{2+} concentration, fractional Ca^{2+} release, and resting SR Ca^{2+} concentration were recorded in FDB fibers from young adult mice using the synthetic Ca^{2+} indicator Fluo-5N [43]. In contrast with the results reported here, the authors reported 88% SR Ca^{2+} depletion in response to 10 s/50 Hz tetanic stimulation. Although they used long pulses, the reported level of depletion is larger than that recorded with D1ER or CatchER; luminal Ca^{2+} depletion seems to plateau when measured with these biosensors, possibly due to an

equilibrium in SR Ca^{2+} release and uptake. The Fluo-5N SR-loading method allowed the first quantitative estimates of static content and dynamic changes in SR Ca^{2+} concentration, but the prolonged dye-loading process limits its use in physiological recordings.

We demonstrated previously that resting SR Ca^{2+} concentration does not differ in fibers from young and old mice; however, SR residual Ca^{2+} was not quantified in old mice. Here, for the first time, we report that residual Ca^{2+} increases with aging, which fully supports our theory that SR Ca^{2+} availability does not limit SR Ca^{2+} release.

Residual SR Ca^{2+} after prolonged myofiber depolarization

We demonstrated previously using confocal microscopy that CatchER expresses in the myofiber SR [34]. Here, we show that blocking RyR1 results in complete ablation of CatchER's fluorescence signal. We rule out the possibility that CatchER is located in the cytosol because, in addition to the morphological evidence, basal fluorescence is intense, which indicates that the biosensor is exposed to high Ca^{2+} concentration. Thus, exposed to high Ca^{2+} concentrations, CatchER signal decreases in response to sarcolemmal depolarization. In contrast, the cytosolic compartment of FDB fibers is strongly buffered, and basal free Ca^{2+} is below 100 nM [12], preventing significant CatchER fluorescence under resting conditions.

Whether SR Ca^{2+} recovery in response to fiber excitation fully represents SR Ca^{2+} uptake from the cytosol, or the signal is influenced by Ca^{2+} redistribution between bound and unbound conformations of the biosensor, calsequestrin's buffer capacity, or Ca^{2+} diffusion from the longitudinal to the terminal cisternae SR is not known. CatchER measurements in a calsequestrin knockout mouse model together with a more refined optical detector, such as a two-photon or a spot scanning confocal microscope [5], in stretched muscle fibers might answer these questions. Previous studies assessed SR Ca^{2+} in amphibian and mammalian skeletal and cardiac muscle SR [12, 14, 23, 28, 29]. The use ofameleon Ca^{2+} sensors represented an advance over previous methods. First, they can be selectively targeted to intracellular organelles in intact cells or tissues in vivo or in vitro. Second, their ratiometric property minimizes confounding factors, such as mechanical artifacts and probe concentration.

In this work, we found a significant residual SR Ca^{2+} after prolonged myofiber depolarization, and we recorded a larger SR Ca^{2+} depletion in myofibers from young compared to old mice. Instead of single pulse, we used a more physiological repetitive stimulation. A train of pulses was sustained until CatchER's signal reached a plateau. Under these experimental conditions, a more significant residual Ca^{2+} was recorded in the SR from old mice. Residual SR Ca^{2+} was quantified based on converting CatchER's signal

into Ca^{2+} concentration according to published methods [34, 43]. These experiments support the conclusion that CatchER is suitable for measuring Ca^{2+} in a high concentration cellular environment; that residual SR Ca^{2+} concentration is significant beyond physiological demands; and that it is higher in myofibers from old compared to young adult mice. CatchER's relatively low affinity measured in mouse FDB myofibers [34] makes it appropriate for detecting SR Ca^{2+} concentration and dynamics. The myofiber sustains SR release in response to repetitive stimulation probably due to efficient SR Ca^{2+} pump-mediated Ca^{2+} uptake. Differences in residual SR Ca^{2+} in response to sarcolemmal depolarization in myofibers from young and old mice reflect excitation–contraction uncoupling, the molecular substrate of which is the age-dependent decline in Cav1.1 expression and a larger number of uncoupled RyR1 [25, 40]. We propose that releasing a larger residual Ca^{2+} pool in myofibers by increasing Cav1.1 expression will improve myofiber-specific force and power with aging.

Acknowledgments The present study was supported by grants from the National Institutes of Health/National Institute on Aging AG07157, AG33385, and AG15820 to Osvaldo Delbono; and EB007268 and GM081749 to Jenny Yang, and the Wake Forest University Claude D. Pepper Older Americans Independence Center (P30-AG21332).

References

- Canato M, Scorsetto M, Giacomello M, Protasi F, Reggiani C, Stienen GJ (2010) Massive alterations of sarcoplasmic reticulum free calcium in skeletal muscle fibers lacking calsequestrin revealed by a genetically encoded probe. *Proc Natl Acad Sci U S A* 107:22326–22331
- Conti A, Gorza L, Sorrentino V (1996) Differential distribution of ryanodine receptor type 3 (RyR3) gene product in mammalian skeletal muscles. *Biochem J* 316(Pt 1):19–23
- Delbono O (2011) Excitation–contraction coupling regulation in aging skeletal muscle. In: Lynch GS (ed) *Sarcopenia—age-related muscle wasting and weakness. Mechanisms and treatment*. Springer, Victoria, pp 113–134
- Demaurex N, Frieden M (2003) Measurements of the free luminal ER Ca^{2+} concentration with targeted “cameleon” fluorescent proteins. *Cell Calcium* 34:109–119
- DiFranco M, Quinonez M, DiGregorio DA, Kim AM, Pacheco R, Vergara JL (1999) Inverted double-gap isolation chamber for high-resolution calcium fluorimetry in skeletal muscle fibers. *Pflügers Arch* 438:412–418
- DiFranco M, Neco P, Capote J, Meera P, Vergara JL (2006) Quantitative evaluation of mammalian skeletal muscle as a heterologous protein expression system. *Protein Expr Purif* 47:281–288
- Fryer MW, Stephenson DG (1996) Total and sarcoplasmic reticulum calcium contents of skinned fibres from rat skeletal muscle. *J Physiol* 493(Pt 2):357–370
- González A, Ríos E (1993) Perchlorate enhances transmission in skeletal muscle excitation–contraction coupling. *J Gen Physiol* 102(3):373–421
- Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ (1981) Improved patch-clamp techniques for high-resolution current recording from cells and cell-free patches. *Pflügers Arch* 391:85–100
- Hidalgo C, Donoso P, Rodriguez PH (1996) Protons induce calsequestrin conformational changes. *Biophys J* 71:2130–2137
- Jimenez-Moreno R, Wang ZM, Gerring R, Delbono O (2008) Sarcoplasmic reticulum Ca^{2+} release declines in muscle fibers from aging mice. *Biophys J* 94:3178–3188
- Jimenez-Moreno R, Wang ZM, Messi ML, Delbono O (2010) Sarcoplasmic reticulum Ca^{2+} depletion in adult skeletal muscle fibres measured with the biosensor D1ER. *Pflügers Arch* 459:725–735
- Jones LM, Yang W, Maniccia AW, Harrison A, van der Merwe PA, Yang JJ (2008) Rational design of a novel calcium-binding site adjacent to the ligand-binding site on CD2 increases its CD48 affinity. *Protein Sci* 17:439–449
- Kabbara AA, Allen DG (1999) Measurement of sarcoplasmic reticulum Ca^{2+} content in intact amphibian skeletal muscle fibres with 4-chloro-m-cresol. *Cell Calcium* 25:227–235
- Lewis RS (2007) The molecular choreography of a store-operated calcium channel. *Nature* 446:284–287
- Melzer W, Herrmann-Frank A, Lüttgau HC (1995) The role of Ca^{2+} ions in excitation–contraction coupling of skeletal muscle fibres. *Biochim Biophys Acta* 1241:59–116
- Owen VJ, Lamb GD, Stephenson DG, Fryer MW (1997) Relationship between depolarization-induced force responses and Ca^{2+} content in skeletal muscle fibres of rat and toad. *J Physiol* 498(Pt 3):571–586
- Palmer AE, Jin C, Reed JC, Tsien RY (2004) Bcl-2-mediated alterations in endoplasmic reticulum Ca^{2+} analyzed with an improved genetically encoded fluorescent sensor. *Proc Natl Acad Sci U S A* 101:17404–17409
- Palmer AE, Giacomello M, Kortemme T, Hires SA, Lev-Ram V, Baker D, Tsien RY (2006) Ca^{2+} indicators based on computationally redesigned calmodulin-peptide pairs. *Chem Biol* 13:521–530
- Pape PC, Jong DS, Chandler WK (1995) Calcium release and its voltage dependence in frog cut muscle fibers equilibrated with 20 mM EGTA. *J Gen Physiol* 106:259–336
- Park KS, Poburko D, Wollheim CB, Demareux N (2009) Amiloride derivatives induce apoptosis by depleting ER Ca^{2+} stores in vascular endothelial cells. *Br J Pharmacol* 156:1296–1304
- Payne AM, Zheng Z, Gonzalez E, Wang ZM, Messi ML, Delbono O (2004) External Ca^{2+} -dependent excitation–contraction coupling in a population of ageing mouse skeletal muscle fibres. *J Physiol* 560(1):137–157
- Picht E, DeSantiago J, Blatter LA, Bers DM (2006) Cardiac alternans do not rely on diastolic sarcoplasmic reticulum calcium content fluctuations. *Circ Res* 99:740–748
- Posterino GS, Lamb GD (2003) Effect of sarcoplasmic reticulum Ca^{2+} content on action potential-induced Ca^{2+} release in rat skeletal muscle fibres. *J Physiol* 551:219–237
- Renganathan M, Messi ML, Delbono O (1997) Dihydropyridine receptor-ryanodine receptor uncoupling in aged skeletal muscle. *J Membr Biol* 157:247–253
- Rossi D, Sorrentino V (2002) Molecular genetics of ryanodine receptors Ca^{2+} -release channels. *Cell Calcium* 32:307–319
- Rudolf R, Mongillo M, Rizzuto R, Pozzan T (2003) Looking forward to seeing calcium. *Nat Rev Mol Cell Biol* 4:579–586
- Rudolf R, Magalhaes PJ, Pozzan T (2006) Direct in vivo monitoring of sarcoplasmic reticulum Ca^{2+} and cytosolic cAMP dynamics in mouse skeletal muscle. *J Cell Biol* 173:187–193
- Shannon TR, Pogwizd SM, Bers DM (2003) Elevated sarcoplasmic reticulum Ca^{2+} leak in intact ventricular myocytes from rabbits in heart failure. *Circ Res* 93:592–594
- Shirokova N, Garcia J, Pizarro G, Ríos E (1996) Ca^{2+} release from the sarcoplasmic reticulum compared in amphibian and mammalian skeletal muscle. *J Gen Physiol* 107:1–18
- Somlyo AV, Gonzalez-Serratos HG, Shuman H, McClellan G, Somlyo AP (1981) Calcium release and ionic changes in the

- sarcoplasmic reticulum of tetanized muscle: an electron-probe study. *J Cell Biol* 90:577–594
32. Sztretye M, Yi J, Figueroa L, Zhou J, Royer L, Allen P, Brum G, Rios E (2011) Measurement of RyR permeability reveals a role of calsequestrin in termination of SR Ca^{2+} release in skeletal muscle. *J Gen Physiol* 138:231–247
 33. Sztretye M, Yi J, Figueroa L, Zhou J, Royer L, Rios E (2011) D4cpv-calsequestrin: a sensitive ratiometric biosensor accurately targeted to the calcium store of skeletal muscle. *J Gen Physiol* 138:211–229
 34. Tang S, Wong H-C, Wang ZM, Huang Y, Zou J, Zhuo Y, Pennati A, Gadda G, Delbono O, Yang JJ (2011) Design and application of a class of sensors to monitor Ca^{2+} dynamics in high Ca^{2+} concentration cellular compartments. *Proc Natl Acad Sci* 13:16265–16270
 35. Taylor JR, Zheng Z, Wang ZM, Payne AM, Messi ML, Delbono O (2009) Increased CaVbeta1A expression with aging contributes to skeletal muscle weakness. *Aging Cell* 8:584–594
 36. Treves S, Vukcevic M, Maj M, Thurnheer R, Mosca B, Zorzato F (2009) Minor sarcoplasmic reticulum membrane components that modulate excitation–contraction coupling in striated muscles. *J Physiol* 587:3071–3079
 37. Ursu D, Schuhmeier RP, Melzer W (2005) Voltage-controlled Ca^{2+} release and entry flux in isolated adult muscle fibres of the mouse. *J Physiol* 562:347–365
 38. Volpe P, Simon BJ (1991) The bulk of Ca^{2+} released to the myoplasm is free in the sarcoplasmic reticulum and does not unbind from calsequestrin. *FEBS Lett* 278:274–278
 39. Wang ZM, Messi ML, Delbono O (1999) Patch-clamp recording of charge movement, Ca^{2+} current and Ca^{2+} transients in adult skeletal muscle fibers. *Biophys J* 77:2709–2716
 40. Wang Z-M, Messi ML, Delbono O (2000) L-type Ca^{2+} channel charge movement and intracellular Ca^{2+} in skeletal muscle fibers from aging mice. *Biophys J* 78:1947–1954
 41. Woods CE, Novo D, DiFranco M, Capote J, Vergara JL (2005) Propagation in the transverse tubular system and voltage dependence of calcium release in normal and mdx mouse muscle fibres. *J Physiol* 568:867–880
 42. Xi Q, Adebiyi A, Zhao G, Chapman KE, Waters CM, Hassid A, Jaggar JH (2008) IP3 constricts cerebral arteries via IP3 receptor-mediated TRPC3 channel activation and independently of sarcoplasmic reticulum Ca^{2+} release. *Circ Res* 102:1118–1126
 43. Ziman AP, Ward CW, Rodney GG, Lederer WJ, Bloch RJ (2010) Quantitative measurement of Ca^{2+} in the sarcoplasmic reticulum lumen of mammalian skeletal muscle. *Biophys J* 99:2705–2714
 44. Zou J, Hofer AM, Lurtz MM, Gadda G, Ellis AL, Chen N, Huang Y, Holder A, Ye Y, Louis CF, Welshhans K, Rehder V, Yang JJ (2007) Developing sensors for real-time measurement of high Ca^{2+} concentrations. *Biochemistry* 46:12275–12288