



Optical spectroscopy studies of the interaction between thiophanate methyl and human serum albumin for biosensor applications

I.S. Yahia^{a,c}, Abdulaziz A. Al-Khedhairi^b, Javed Musarrat^{b,c}, F. Yakuphanoglu^{d,*}

^a Department of Physics, Faculty of Education, Ain Shams University, Roxy, Cairo, Egypt

^b Al-Jeraisy Chair for DNA Research, Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^c Department of Microbiology, Faculty of Agricultural Sciences, Aligarh Muslim University, Aligarh, India

^d Department of Metallurgical and Materials Engineering, Firat University, Elazığ, Turkey

ARTICLE INFO

Article history:

Received 15 December 2010

Received in revised form 14 April 2011

Accepted 27 April 2011

Keywords:

Biosensor

Human serum albumin (HSA)

Thiophanate methyl (MT)

Optical constants

AFM

ABSTRACT

Optical properties of the interaction between thiophanate methyl and human serum albumin have been investigated for biosensor applications. The interaction between human serum albumin (HSA) and thiophanate methyl (MT) was investigated by UV–Vis absorption spectra and atomic force microscopy. The optical constants (refractive index, absorption index, band gap and dielectric properties) of HSA, MT and MT + HSA films were determined using absorbance, transmittance and reflectance spectra. The refractive index dispersion curve (>530 nm) exhibits the normal dispersion. The refractive index of the MT + HSA is higher than both HSA and MT alone due to the highest reflectance of the mixture of MT and HSA. This behavior is indicative of the complex formation between the MT and HSA.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Human serum albumin (HSA), represents the major soluble protein constituent of the circulatory system. It is the most abundant drug carrier protein which plays an important role in the transportation and deposition of many endogenous and exogenous drugs/ligands in blood [1,2]. Many drugs, chemicals and bioactive small molecules can bind reversibly to albumin and other serum components, which implicate their role as carriers [3–5]. HSA can bind and carry many drugs and chemical materials which are soluble or poorly soluble in water. The distribution, free concentration and the metabolism of various drugs can be obviously altered upon binding with HSA [1]. Modern agricultural production in many countries largely depends on intensive use of pesticides. Indiscriminate applications of pesticides resulted in accumulation of pesticide residues in fruits and vegetables [6–9]. Thiophanate methyl (MT) is a thioallophanate compound, which is widely used to control the fungal diseases of crops due to its broad spectrum activity [10,11]. Apart from the beneficial effects associated with the use of pesticides, the pesticides may also pose potential hazards to humans and the environment [12,13]. MT is a known category-III acute inhalation toxicant, likely to be carcinogenic to

humans and is widely used for the control of some important fungal diseases of crop [14–16]. It is a potential spindle poison and impairs the polymerization of tubulin [17] besides causing delayed cellular proliferation and increase in the frequency of apoptosis. Recently, Saquib et al. [18,19] demonstrated its genotoxic potential and capacity to induce DNA strand breaks, micronuclei and 8-oxodG formation in human lymphocytes due to ROS generation in vitro. The interactions between bio-macromolecules and chemicals have attracted increasing research interest in recent years especially for the health and environmental protection [20,21]. Among various bio-macromolecules, serum albumins are the most abundant soluble proteins in the body circulatory system of a wide variety of organisms and have many indispensable physiological functions. Serum albumins are serving as a depot protein and a transport protein for many exogenous compounds [14,21]. Therefore, the absorption, distribution, metabolism and excretion properties, as well as the stability and toxicity of chemical substances can be significantly affected because of their binding to serum albumins [14–21]. Moreover, there is evidence of conformational changes of serum albumin induced by its interaction with low molecular weight molecules, which appear to affect the secondary and tertiary structure of albumins [14–21]. Therefore, the studies on interaction of chemicals with serum albumin are of great importance. Binding of chemicals with protein helps in understanding the metabolism and transport process besides assessing the structure–function relationship of the protein [20]. Thus, it is necessary to investigate the interaction of MT with HSA, to

* Corresponding author. Tel.: +90 5337623815.

E-mail addresses: fyhanoglu@firat.edu.tr, fyhan@hotmail.com (F. Yakuphanoglu).

demonstrate the MT–protein complexation and provide important insight into the nature of interaction. The interaction has been studied by means of spectrophotometric studies and AFM. UV–Vis–NIR spectra of transmittance/reflectance/absorbance were measured in the wavelength range 250–800 nm. Optical constants of HSA, MT and complex interaction between them are calculated and analyzed in the light of the optical spectroscopy for the first time to characterize the interaction between MT and HSA.

AFM micrographs as a powerful tool provide more useful information about the complex interaction between MT and HSA.

2. Experimental procedures

2.1. Chemicals and materials

HSA was purchased from Sigma–Aldrich and MT was obtained from D-86199 Augsburg (Germany).

2.2. Reagents

HSA solution was prepared in pH 7.4 buffer solution as (3.0×10^{-5} mol/L) and HSA stock solutions were kept in the dark at room temperature (293 K). The stock solution (1.0×10^{-3} mol/L) of MT was prepared in ethanol.

2.3. Instrumentation

Thin film of HSA and MT and MT + HSA were prepared through Xe-100 Desk-top Spin Coater at 1000 rpm and dried at 60 °C for 5 min. The transmittance/reflectance/absorbance spectra of the HSA, MT and MT + HSA thin films were obtained using a Shimadzu UV-VIS-NIR 3600 spectrophotometer. The morphology of MT and MT + HSA thin films was investigated using Park System XE-100E atomic force microscope.

3. Results and discussion

3.1. AFM micrographs of MT and MT + HSA thin films

The atomic force microscopy (AFM) has been exploited as a surface characterization technique for examination of biomolecules in biomaterial research. This technique offers the significant advantage of probing in high detail the surface topography qualitatively (surface images) and quantitatively (surface roughness) due to its nanometer-scale spatial resolution. It has proved to be very helpful for the determination and verification of various morphological features and parameters. The interaction between HSA and MT in thin film form was examined. Two and three dimensional AFM micrographs of $40 \times 40 \mu\text{m}^2$ and $10 \times 10 \mu\text{m}^2$ of MT and MT + HSA are shown in Fig. 2. The MT thin film micrographs dried at 60 °C have the shape of tree leaves with roughness 85.977 nm. The mixed MT and HSA showed the roughness of 75.369 nm, suggestive of strong binding energy between MT and HSA. Li et al. [6] and also reported the interaction between MT and HSA via the UV absorption spectra, the fluorescence emission and Fourier transform infrared (FT-IR) spectroscopy and the molecular modeling studies. The effect of MT on HSA caused considerable changes in the protein secondary structure, and the AFM data support the interaction between MT with human serum protein. These results corroborate with our earlier studies on circular dichroism (CD) analysis [22], which have explicitly demonstrated the MT-induced conformational changes in the secondary structure of protein. The significant reductions in the mean residue ellipticity (MRE) in $\text{deg cm}^2 \text{dmol}^{-1}$, suggested alterations in protein helicity upon MT–HSA complexation. The MT concentration dependent decrease in band intensities at 209 and




Fig. 1. The chemical structure of MT.

222 nm in the far-UV region reflected a change in the protein secondary structure, primarily due to reduction in α -helical content of treated protein. At a molar ratio of 1:0.04, the perturbation has been determined to be 8% with loss in protein helicity from 54.8% in native untreated HSA to 50.6% upon MT treatment [22].

3.2. Optical Transmittance/reflectance/absorbance of HSA, MT and MT + HSA

Optical absorbance/transmittance/reflectance spectra were used to determine the optical constants (refractive index, absorption index, band gap and dielectric properties) of HSA, MT and MT + HSA for optical biosensor applications. Both the absorbance and the reflectance were recorded on UV–Vis–NIR double beam spectrophotometer at room temperature in the range 250–800 nm. The transmittance/reflectance spectra of HSA, MT and MT + HSA are shown in Fig. 3a. The data revealed that the HSA, MT and MT–HSA complex exhibit a transparent region through a wide range of wavelengths as shown in Fig. 1a. It is clear that the transmittance of a mixture (MT + HSA) is lower than the MT and HSA alone, indicating that the interaction occurred between them. The reflectance of MT + HSA was significantly higher than MT and HSA alone, which has affected the calculated optical constants of MT, HSA and the mixture (MT + HSA). UV–Vis absorption measurement is a very simple but effective method to explore the structural change and the complex formation. Fig. 4b represents the UV–Vis–NIR absorption spectra of HSA, MT and MT–HSA complex binding in which the most absorption area of this materials are in UV region as shown in the inset of this figure. Also, the HSA exhibited higher absorbance than both MT and MT + HSA. It is also found that the complex interaction between HSA and MT led to the enhancement the absorbance to be increased than MT alone and lower than HSA spectra. Accordingly, the MT–HSA complex formation upon mixing MT with HSA results in the change in protein conformation due to change in the micro-environment around HSA. The results correspond well with our earlier studies, which have suggested that the MT–HSA complex formation occurs mainly due to binding with aromatic amino acids, preferably the tryptophan moiety on sub-domain IIA due to the alkyl group transfer from MT to amino acid functional group [22]. The complex formation has also been validated by cyclic voltammetric analysis, where a significant reduction in the cathodic peak current (ipc) was noticed upon addition of HSA to MT solution. The reduction in peak current is attributed to the HSA interaction with MT to form a non-electrochemical complex, which blocks the electron transfer between the quasi-reversible peaks of MT and electrode [22].

Refractive index measurement plays a vital role in many areas of biophysics, biochemistry and biomedicine. The complex refractive index and dielectric function characterize the optical properties of thin materials. The optical properties of the thin films have been investigated by spectrophotometric measurement of transmittance/reflectance/absorbance at the wavelengths range of 250–800 nm. The refractive index was calculated through the following equations. The normal reflectance




Fig. 2. AFM images of (2D 40 × 40 μm²) and (2D 10 × 10 μm²) of MT and MT + HSA. Insets: three dimensional (3D) AFM images (3D 40 × 40 μm²) and (3D 10 × 10 μm²) for MT and MT + HSA.

for any absorbing material can be expressed through the following equation [23–25]:

$$R = \left(\frac{(n-1)^2 + k^2}{(n+1)^2 + k^2} \right). \quad (1)$$

Solution of Eq. (1) gives directly the refractive index as [24,26,27]:

$$n = \left(\frac{1+R}{1-R} \right) + \sqrt{\left(\frac{4R}{1-R} - k^2 \right)}, \quad (2)$$

where n is the refractive index and k is absorption index. The values of the absorption index k can be calculated from the absorption coefficient $\alpha = A/t$, where A is the measured absorbance and t the thickness of the investigated films measured through AFM. The variation of refractive index (n) and extinction coefficient (k) with wavelength for HSA, MT and the MT–HSA complex is shown in Fig. 4. It is observed that the refractive index possesses a minimum and maximum peak (anomalous dispersion) at the strong absorption region while at the higher wavelengths, all the samples exhibit a normal dispersion behavior in which the refractive index is decreased with increasing wavelengths. Also, the refractive index of the MT + HSA is higher than both HSA and MT due to the highest reflectance of the mixture (MT and HSA). This behavior is an indicator of the complex formation between the MT and HSA. For the absorption index, it is clear that HSA has the strongest absorption than both MT and MT + HSA. Also, MT + HSA exhibits the absorption

between HSA and MT. The previous work on HSA shows that HSA exhibits a strong absorption in the UV region [28].

3.2.1. Determination of band gaps

The fundamental absorption is related to band-to-band or to exciton transition. The relation between absorption coefficient and optic band gap for direct and indirect transitions is given by the following equation [29,30]:

$$\alpha h\nu = A(h\nu - E_g)^r, \quad (3)$$

where α is the absorbance coefficient, $h\nu$ is the photon energy, E_g is the optical band gap, A is a constant which does not depend on photon energy and r has four numeric values (1/2 for allowed direct, 2 for allowed indirect, 3 for forbidden direct and 3/2 for forbidden indirect optical transitions). In this work, direct band gap is determined by plotting $(\alpha h\nu)^2$ versus $h\nu$ curve. With the extrapolation of the linear region to lower energies, it is possible to calculate the band gap of HSA, MT and the MT + HSA. The optical band gaps of HSA, MT and the MT + HSA were found to be 4.10, 3.98 and 3.92 eV, respectively. The optical band gap of MT changes with interaction between MT and HSA.

For optical materials, another important parameter is the refractive index. The complex refractive index can be described by [25,31,32]:

$$\hat{n} = n(h\nu) + ik(h\nu). \quad (4)$$




Fig. 3. (a and b) Transmittance/reflectance spectra of HSA, MT and MT + HSA compounds.

Also, the complex dielectric constant function can be written as follows:

$$\hat{\epsilon} = \epsilon_1(h\nu) + i\epsilon_2(h\nu). \quad (5)$$

Both the complex refractive index and the dielectric constant characterize the optical properties of thin film materials. Where $\epsilon_1(h\nu)$




Fig. 4. The UV absorbance of HSA, MT and MT + HSA compounds.




Fig. 5. (a and b) The variation of refractive index and absorption index with wavelength for HSA, MT and MT + HSA compounds.

and $\epsilon_2(h\nu)$ is the real and imaginary parts of the dielectric constant. Both $\epsilon_1(h\nu)$ and $\epsilon_2(h\nu)$ are related to the refractive index and absorption index by the following equations [31,32]:

$$\epsilon_1(h\nu) = n^2(h\nu) - k^2(h\nu) \quad (6)$$

and

$$\epsilon_2(h\nu) = 2n(h\nu) \cdot k(h\nu). \quad (7)$$

The dependence of both $\epsilon_1(h\nu)$ and $\epsilon_2(h\nu)$ on photon energy is shown in Fig. 5(a and b), respectively. Both $\epsilon_1(h\nu)$ and $\epsilon_2(h\nu)$




Fig. 6. Plot of $(\alpha h\nu)^2$ versus the photon energy $h\nu$ for the HSA, MT and MT + HSA compounds.




Fig. 7. (a and b) Plots of $\epsilon_1(h\nu)$ and $\epsilon_2(h\nu)$ of HSA, MT and MT + HSA compounds.

increase with increasing the photon energy until reached its maximum and then they are decreased with the further increase of the photon energy. It is clear that, the values of $\epsilon_1(h\nu)$ is larger than of $\epsilon_2(h\nu)$. Also, the values of $\epsilon_1(h\nu)$ is larger for MT+HSA than MT and HSA alone, due to the high refractive index of the mixture of MT+HSA than MT and HSA alone. For the imaginary part of dielectric constant, the value for HSA is larger than MT and the mixture MT + HSA is in the middle between them followed the same pattern of the absorbance spectrum and absorption index (see Figs. 6–8).




Fig. 8. Plots of the dielectric relaxation time τ versus the photon energy $h\nu$ for HSA, MT and MT + HSA compounds.

The dielectric relaxation time τ can be determined by the following equation [33,34]:

$$\tau = \frac{\epsilon_\infty - \epsilon_1}{\omega \epsilon_2} \quad (9)$$

The dielectric relaxation time τ as a function of photon energy for HSA, MT and MT + HSA films. This figure shows that the relaxation time is decreased with increasing photon energy. Also, the values of MT + HSA at the lower photon energy are much higher than MT and HSA alone. Such behavior was expected due to the interaction between them forming a complex of MT + HSA.

4. Conclusions

The interaction between thiophanate methyl and human serum albumin has been investigated by UV–Vis absorption spectra and atomic force microscopy. The optical constants such as refractive index, absorption index, band gap and dielectric properties of HSA, MT and MT + HSA films were determined. The optical constants of MT change with HSA interaction. This suggests that the optical biosensors can be prepared by MT + HSA films.

Acknowledgments

This work was supported by Global Research Network for Electronic Devices & Biosensors (GRNEDB) and KING Saud University, Riyadh, Saudi Arabia.

References

- [1] J. Li, C. Ren, Y. Zhang, X. Liu, X. Yao, Z. Hu, J. Mol. Struct. 881 (2008) 90–96.
- [2] H. Vorum, B. Honore, J. Pharm. Pharmacol. 48 (1996) 870.
- [3] D.C. Carter, B. Chang, J.X. Ho, K. Keeling, Z. Krishnasami, Eur. J. Biochem. 226 (1994) 1049.
- [4] J. Guharay, B. Sengupta, P.K. Sengupta, Proteins 43 (2001) 75.
- [5] Y. Li, W.Y. He, J.Q. Liu, F.L. Sheng, Z.D. Hu, X.G. Chen, Gen. Subj. 1722 (2005) 15.
- [6] J. Li, X. Liu, C. Ren, J. Li, F. Sheng, Z. Hu, J. Photochem. Photobiol. B 94 (2009) 158.
- [7] F.V. Bright, M.E.P. McNally, Supercritical fluid technology: theoretical and applied approaches to analytical chemistry, in: ACS Symp. Ser., vol. 488, American Chemical Society, Washington, DC, 1992.
- [8] C.M. Torres, Y. Picó, J. Maries, J. Chromatogr. A 754 (1996) 301.
- [9] C. Blasco, G. Font, Y. Picó, Food Control 17 (2006) 841.
- [10] J.H. Canton, Bull. Environ. Contam. Toxicol. 16 (1976) 214.
- [11] M.E. Traina, P. Fazzi, C. Macri, C. Ricciardi, A.V. Stazi, E. Urbani, A. Mantovani, J. Appl. Toxicol. 18 (1998) 241.
- [12] WHO, The WHO Recommended Classification of Pesticides by Hazard and Guidelines to the Classification 1988–1989, World Health Organisation, Geneva, 1988.
- [13] D.J. Ecobichon, Toxic effects of pesticides, in: C.D. Klaassen (Ed.), Casarett and Doull's Toxicology, The Basic Science of Poisons, McGraw-Hill, New York, 2001, pp. 763–810; in: Proposed Guidelines for Carcinogen Risk Assessment, Office of Research and Development U.S. Environmental Protection Agency Washington, DC, EPA/600/P-92/003C, Fed. Regist. 61 (79) (1996) 17960; K.A. Hassall, The Biochemistry and Uses of Pesticides, second ed., Macmillan Press Ltd., 1990; M.E. Traina, P. Fazzi, C. Macri, C. Ricciardi, A.V. Stazi, E. Urbani, A. Mantovani, J. Appl. Toxicol. 18 (1998) 241; F. Maranghi, C. Macri, C. Ricciardi, A.V. Stazi, M. Resci, A. Mantovani, Reprod. Toxicol. 17 (2003) 617.
- [14] M. Paolini, L. Pozzetti, P. Perocco, M. Mazzullo, G. Cantelli-Forti, Cancer Lett. 135 (1999) 203.
- [15] N. Shahabadi, M. Maghsudi, Z. Kiani, M. Pourfoulad, Food Chem. 124 (2011) 1063.
- [16] Y. Li, W. He, J. Liu, F. Sheng, Z. Hu, X. Chen, Biochem. Biophys. Acta 1722 (2005) 15.
- [17] H.X. Sun, Y.L. Tang, J.F. Xiang, G.Z. Xu, Y.Z. Zhang, H. Zhang, et al., Bioorg. Med. Chem. Lett. 16 (2006) 3586.
- [18] Q. Saquib, A.A. Al-Khedhairi, S. Al-Arif, A. Dhawan, J. Musarrat, Toxicol. In Vitro 23 (2009) 848.
- [19] Q. Saquib, A.A. Al-Khedhairi, B.R. Singh, J.M. Arif, J. Musarrat, J. Environ. Sci. Health (B) 45 (2010) 1.
- [20] J. Flarakos, K.L. Morand, P. Vouros, Anal. Chem. 77 (2005) 1345.
- [21] T.O. Hushcha, A.I. Luik, Y.N. Naboka, Talanta 53 (2005) 29.

- [22] Q. Saquib, A.A. Al-Khedhairy, S. Al-Arifi, S. Dwivedi, J. Mustafa, J. Musarrat, Int. J. Biol. Macromol. 47 (2010) 60.
- [23] M. Caglar, S. Ilican, Y. Caglar, F. Yakuphanoglu, Appl. Surf. Sci. 255 (2009) 4491.
- [24] F. Abeles (Ed.), Optical Properties of Solids, North-Holland, Publishing Company, London, UK, 1972.
- [25] J.N. Hodgson, Optical Absorption and Dispersion in Solids, Chapman and Hall Ltd., 11 New Fetter Lane, London EC4, 1970.
- [26] F. Yakuphanoglu, G. Barım, I. Erol, Physica B 391 (2007) 136.
- [27] F. Yakuphanoglu, M. Kandaz, M.N. Yaraşır, F.B. Şenkal, Physica B 393 (2007) 235.
- [28] L. Trnková, I. Boušová, V. Staňková, J. Dršata, J. Mol. Struct. 985 (2011) 243–250.
- [29] O.A. Azim, M.M. Abdel-Aziz, I.S. Yahia, Appl. Surf. Sci. 255 (2009) 4829.
- [30] F. Yakuphanoglu, J. Alloys Compd. 470 (2009) 55.
- [31] A.A.M. Farag, I.S. Yahia, Opt. Commun. 283 (2010) 4310.
- [32] F. Yakuphanoglu, M. Sekerci, O.F. Ozturk, Opt. Commun. 239 (2004) 275.
- [33] M.M. Abdel-Aziz, I.S. Yahia, L.A. Wahab, M. Fadel, M.A. Afifi, Appl. Surf. Sci. 252 (2006) 8163.
- [34] E.G. El-Metwally, M.O. Abou-Helal, I.S. Yahia, J. Ovon. Res. 4 (2008) 20.