

# Bacterial Whole-Cell Biosensor for Glutamine with Applications for Quantifying and Visualizing Glutamine in Plants

Michael J. Tessaro,<sup>a</sup> Sameh S. M. Soliman,<sup>a,b</sup> and Manish N. Raizada<sup>a</sup>

Department of Plant Agriculture, University of Guelph, Guelph, Ontario, Canada,<sup>a</sup> and Department of Pharmacognosy, Faculty of Pharmacy, Zagazig University, Zagazig, Egypt<sup>b</sup>

**A whole-cell biosensor for glutamine (GlnLux) was constructed by transforming an *Escherichia coli* glutamine (Gln) auxotroph with a constitutive *lux* reporter gene. Measurements of Gln in plant extracts using GlnLux correlated with quantification using high-performance liquid chromatography (Spearman's  $r = 0.95$ ). GlnLux permitted charge-coupled-device (CCD) imaging of Gln from whole plant organs.**

Microbial whole-cell biosensors have been used to quantify analytes from biological extracts into which they are coinoculated (6, 8, 21). Such biosensors have also been employed for *in vivo* imaging of metabolites following their leakage from plant organs onto agar embedded with biosensor bacteria expressing a visible reporter gene (12, 14, 19, 22).

Glutamine (Gln) is a critical intermediate in nitrogen metabolism. In plants, soil nitrogen is assimilated into Gln, which then serves as a nitrogen donor for other biochemical reactions (18). Gln is also a major long-distance transport form of nitrogen (e.g., root to shoot) (20). Today, free Gln is primarily quantified using tissue extracts by high-performance liquid chromatography (HPLC) (1, 17), not *in vivo* imaging.

Here we constructed a microbial *lux* biosensor for Gln. *Escherichia coli* Gln-auxotrophic strain JW3841-1 (*E. coli* Genetic Stock Center CGSC 10775, Yale University) was previously generated by disrupting *glnA* with a kanamycin cassette (*glnA732Δ::Kan*) (2). The auxotroph failed to grow in M9 minimal medium except when it was supplemented with Gln. We transformed JW3841-1 with pT7-*lux* (Amp<sup>r</sup>) (16), containing a *luxCDABE* operon from *Vibrio fischeri* driven by a constitutive T7 promoter from *Xenorhabdus luminescens*, creating strain GlnLux.

Gln standards were used to optimize GlnLux for luminometer assays. In the optimized protocol, GlnLux was grown (optical density at 595 nm [OD<sub>595</sub>] of 0.3, LB, 37°C, 250 rpm) and then pelleted (centrifugation at 700 × g, 25°C, 10 min). To deplete endogenous free Gln, the pellet was washed 3 times (in M9 medium lacking Gln), diluted (1/10, OD<sub>595</sub> = 0.025, M9 medium), and then incubated (37°C, 250 rpm, 14 h) (9). Diluting GlnLux cultures in this way resulted in a subsequent reduction of background noise and an increased assay sensitivity (see Fig. S1 in the supplemental material). The culture was then incubated with Gln standards (96-well opaque white plates; Fisher CS003912): GlnLux (10 μl) and M9 medium containing Gln standards (90 μl) were added into each well, covered (sterile breathable film; Sigma A9224), and then incubated (37°C without shaking). Increasing the incubation time of Gln with GlnLux cells shifted the linear range of the assay higher (see Fig. S2 and S3 in the supplemental material). For lux quantification, plates were reread hourly (1-s reads; MicroLumat Plus LB96V; Berthold Technologies, Germany). All media were supplemented with 50 μg/ml kanamycin and 100 μg/ml carbenicillin.

The GlnLux assay using Gln standards achieved a sensitivity of

0.0001 μg/ml (0.68 nM) (see Fig. S2 in the supplemental material), which is similar to that of the most sensitive HPLC-tandem mass spectrometry (HPLC-MS/MS) assay (7, 11). The linear range of the assay extended across 3 orders of magnitude (0.0001 to 0.1 μg/ml Gln) (see Fig. S2 in the supplemental material). GlnLux did not significantly react to other nitrogenous compounds at concentrations expected in plant extracts following a 1,000-fold dilution, the dilution needed to shift extracted Gln into the linear assay range. There was no significant cross-reactivity from NH<sub>4</sub>Cl (≤90 μM, already present in M9 medium), KNO<sub>3</sub> (≤200 μM), NaNO<sub>2</sub> (≤0.3 μM), or a 19-amino-acid cocktail (3 to 110 μM). However, higher nitrate or amino acid concentrations could potentially activate GlnLux (data not shown).

The accuracy of the GlnLux luminometer assay for quantifying Gln in plant extracts was determined by comparing the results with those from HPLC (1) (see the supplemental material for the HPLC methodology and plant treatments). For the GlnLux assay, maize/corn (*Zea mays* L.) leaves were ground (1 to 10 mg) with sand in liquid nitrogen, treated (9 μl double-distilled water [ddH<sub>2</sub>O] with 1% [vol/vol] protease inhibitor cocktail [PIC; Sigma P9599, pH 7.0] per mg), vortexed, and centrifuged (20 min, 4°C, 10,000 × g). The supernatant was transferred to a fresh tube on ice, diluted 1,000-fold (ddH<sub>2</sub>O, pH 7.0), and used immediately. Using 96-well plates, each well was prepared (80 μl 1.25 × M9 medium, 10 μl GlnLux as noted above, 10 μl plant extract). The PIC inhibited suspected proteases in plant extracts (data not shown). An excellent correlation was found between measurements using the GlnLux assay and those using HPLC (Spearman's  $r = 0.95$ ,  $P < 0.0001$ , degrees of freedom [df] = 39) (Fig. 1). Remarkably, GlnLux leaf measurements could distinguish between plants fertilized with and without ammonium/nitrate (Fig. 1) using only the equivalent of 1 μg of tissue per well.

For imaging of Gln in whole plant organs, tissues were freeze-thawed to cause Gln leakage, placed on agar preembedded with

Received 23 April 2011 Accepted 31 October 2011


Published ahead of print 11 November 2011

Address correspondence to Manish N. Raizada, raizada@uoguelph.ca.

Supplemental material for this article may be found at <http://aem.asm.org/>.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.05261-11




**FIG 1** Validation of the GlnLux luminometer assay as a method to quantify free Gln in plant tissue extracts by comparing results to those of a standard HPLC assay. Leaf extracts of maize seedlings previously treated with fertilizer solution with or without nitrogen were quantified using HPLC and independently tested using the GlnLux luminometer assay, with the latter using 1/1,000 dilutions of maize extract, corresponding to 1  $\mu\text{g}$  tissue (fresh weight [FW]). Each value is from an extract from one leaf of one seedling. The coefficient of correlation (Spearman's  $r$ ) between the GlnLux and HPLC assays is indicated. HPLC and GlnLux readings segregated based on the prior nitrogen fertilizer treatment (vertical dotted line), with the exception of two outliers (open circles). The 0- $\mu\text{g}/\text{ml}$  Gln standard reading was subtracted from all lux values, which were read in randomized replicates of 3 to 4 in an endpoint assay set to the "integrate" function. RLU, relative light units.

GlnLux (GlnLux agar), and then imaged using a photon capture camera (22). Briefly, a GlnLux starter culture (LB with 50  $\mu\text{g}/\text{ml}$  kanamycin, 100  $\mu\text{g}/\text{ml}$  carbenicillin, 0.2 mM Gln, 4.0 mM glucose; 16 h, 37°C, 250 rpm) was pelleted (centrifugation at  $1,100 \times g$ , 25°C, 10 to 20 min), washed 3 times (0.01 M potassium phos-

phate buffer, pH 7.0), and resuspended (M9 medium with an  $\text{OD}_{595}$  of 1.0). To prepare GlnLux agar, concentrated M9 medium (with Bacto agar, 10 g/liter) was cooled (42°C), supplemented with Casamino Acids solution (5 g/liter of the final solution), mixed with GlnLux (10% [vol/vol],  $\text{OD}_{595} = 1.0$ ), and then poured. Flash-frozen tissues were thawed (room temperature, 30 s), pressed onto GlnLux agar, inverted, incubated (37°C, 1 to 6 h), and imaged hourly (ChemiProHT, Roper) (4) with a charge-coupled-device (CCD) chip pre-cooled to  $-80^\circ\text{C}$  (5) (200- to 600-s exposures). GlnLux agar permitted imaging of Gln at a sensitivity threshold of 0.01 to 0.1 mM (data not shown).

For plant imaging, maize seedlings were fed with nitrate once or with only water (14 h prior to freeze-thawing). GlnLux agar under nitrate-treated roots emitted a noticeably higher signal than the same agar under water-treated roots (Fig. 2A and B). Leaves were also imaged after their roots were exposed to fertilizer with or without ammonium/nitrate (five times,  $\sim 50$  to 10 h prior to freeze-thawing). The signal from GlnLux agar under leaves from nitrogen-treated seedlings was dramatically higher than that of the same agar under leaves from nitrogen-deficient seedlings (Fig. 2A to H). Thus, GlnLux agar was successful in visualizing root nitrogen uptake, assimilation into Gln, and long-distance transport to leaves (18).

Apart from uptake, free Gln can accumulate at the base of a senescing leaf from scavenged proteins at the tip; scavenged Gln is exported through vascular cells to growing organs (10). In previous studies using maize leaf 2, Gln levels were highest at the leaf base, decreasing by 50% in the middle and by  $>90\%$  near the tip



**FIG 2** Luminescence *in vivo* imaging of Gln in plant organs using GlnLux agar. (A to H) Two-week-old maize seedlings either were treated with nitrogen fertilizer or were not, and harvested plant organs were freeze-thawed to cause Gln leakage and then placed on GlnLux agar; the opposite agar surface was then imaged using a photon capture CCD camera. Representative pictures of a root system not previously exposed to nitrate (A), a root system previously exposed to nitrate (B), leaves from unfertilized plants (C to E), leaves from ammonium nitrate-fertilized plants (F to H), and leaf number 2 from different ammonium nitrate-fertilized plants showing consistent spatial distribution of Gln (I) are shown. (C to H) The leaf tip numbers in each panel from left to right are 1, 2, and 4. The light images (C and F), white lux images (D and G), and false-colored lux images (A, B, E, H, and I) are shown.

(3), consistent with peak glutamine synthetase expression at the leaf base (13, 15). Using GlnLux agar, *lux* expression in fertilized leaf 2 was consistently highest at the base-to-middle zones along the midvein vasculature and lowest at the tip (Fig. 2I). GlnLux agar could thus resolve *in vivo* spatial variation for Gln. Nevertheless, imaging using GlnLux agar is only semiquantitative. Future applications may include high-throughput screening for nitrogen uptake/metabolism mutants. GlnLux is complementary to a recent plant Förster resonance energy transfer (FRET)-based Gln biosensor providing subcellular resolution (23).

## ACKNOWLEDGMENTS

We thank Barry Shelp for HPLC assistance and helpful manuscript comments, Nafisa Tasnim for *lux* visualization assistance, and Mansel Griffiths for pT7-*lux*.

S.S.M.S. received an Egyptian Bureau of Cultural Affairs of Canada Scholarship. M.N.R. was supported by grants from the Ontario Research Fund and OMAFRA.

## REFERENCES

- Allan WL, Shelp BJ. 2006. Fluctuations of gamma-aminobutyrate, gamma-hydroxybutyrate, and related amino acids in *Arabidopsis* leaves as a function of the light-dark cycle, leaf age, and N stress. *Can. J. Bot.* **84**: 1339–1346.
- Baba T, et al. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2**:2006.0008.
- Chapman DJ, Leech RM. 1979. Changes in pool sizes of free amino-acids and amides in leaves and plastids of *Zea mays* during leaf development. *Plant Physiol.* **63**:567–572.
- Chinnusamy V, Stevenson B, Lee B-H, Zhu J-K. 2002. Screening for gene regulation mutants by bioluminescence imaging. *Sci. STKE* **2002**: pl10.
- Christenson MA. 2002. Detection systems optimized for low-light chemiluminescence imaging. In Van Dyke K, Van Dyke C, Woodfork K (ed), *Luminescence biotechnology: instruments and applications*. CRC Press, Boca Raton, FL.
- DeFraia CT, Schmelz EA, Mou ZL. 2008. A rapid biosensor-based method for quantification of free and glucose-conjugated salicylic acid. *Plant Methods* **4**:28.
- Eckstein JA, Ammerman GM, Reveles JM, Ackermann BL. 2008. Analysis of glutamine, glutamate, pyroglutamate, and GABA in cerebrospinal fluid using ion pairing HPLC with positive electrospray LC/MS/MS. *J. Neurosci. Methods* **171**:190–196.
- Erickson AM, Diaz IBZ, Kwon YM, Ricke SC. 2000. A bioluminescent *Escherichia coli* auxotroph for use in an in vitro lysine availability assay. *J. Microbiol. Methods* **40**:207–212.
- Erickson AM, Zabala-Diaz IB, Ricke SC. 1999. Antibiotic amendment for suppression of indigenous microflora in feed sources for an *Escherichia coli* auxotroph lysine assay. *J. Appl. Microbiol.* **87**:125–130.
- Fischer W-N, et al. 1998. Amino acid transport in plants. *Trends Plant Sci.* **3**:188–195.
- Fonteh AN, Harrington RJ, Harrington MG. 2007. Quantification of free amino acids and dipeptides using isotope dilution liquid chromatography and electrospray ionization tandem mass spectrometry. *Amino Acids* **32**: 203–212.
- Jaeger CH, III, Lindow SE, Miller W, Clark E, Firestone MK. 1999. Mapping of sugar and amino acid availability in soil around roots with bacterial sensors of sucrose and tryptophan. *Appl. Environ. Microbiol.* **65**:2685–2690.
- Li PH, et al. 2010. The developmental dynamics of the maize leaf transcriptome. *Nat. Genet.* **42**:1060–1067.
- Loper JE, Lindow SE. 1994. A biological sensor for iron available to bacteria in their habitats on plant surfaces. *Appl. Environ. Microbiol.* **60**: 1934–1941.
- Majeran W, et al. 2010. Structural and metabolic transitions of C4 leaf development and differentiation defined by microscopy and quantitative proteomics in maize. *Plant Cell* **22**:3509–3542.
- Meighen EA, Szittner RB. 1992. Multiple repetitive elements and organization of the *lux* operons of luminescent terrestrial bacteria. *J. Bacteriol.* **174**:5371–5381.
- Micallef BJ, Shelp BJ, Ball RO. 1989. Quantification of C14-labelled amino-acids by reverse-phase high-performance liquid-chromatography. *J. Liq. Chromatogr.* **12**:1281–1300.
- Mifflin BJ, Habash DZ. 2002. The role of glutamine synthetase and glutamate dehydrogenase in nitrogen assimilation and possibilities for improvement in the nitrogen utilization of crops. *J. Exp. Bot.* **53**:979–987.
- Miller WG, Brandl MT, Quinones B, Lindow SE. 2001. Biological sensor for sucrose availability: relative sensitivities of various reporter genes. *Appl. Environ. Microbiol.* **67**:1308–1317.
- Peoples M, Gifford R. 1990. Long-distance transport of carbon and nitrogen from sources to sinks in higher plants, p 442–447. In Dennis DT, Turpin DH (ed), *Plant physiology, biochemistry and molecular biology*. Longman Singapore Publishers, Singapore.
- Pfleger BF, Pitera DJ, Newman JD, Martin VJJ, Keasling JD. 2007. Microbial sensors for small molecules: development of a mevalonate biosensor. *Metab. Eng.* **9**:30–38.
- Soudry E, Ulitzur S, Gepstein S. 2005. Accumulation and remobilization of amino acids during senescence of detached and attached leaves: in planta analysis of tryptophan levels by recombinant luminescent bacteria. *J. Exp. Bot.* **56**:695–702.
- Yang HY, Bogner M, Stierhof YD, Ludewig U. 2010. H<sup>+</sup>-independent glutamine transport in plant root tips. *PLoS One* **5**:e8917.